Agricultu

SEF RELAKE

Agriculture, Food and Natural Resources Journal The Official Journal of the Faculty of Agriculture, Nnamdi Azikkiwe University, Nigeria

ISSN: 3043-5420

Journal homepage: https://journals.unizik.edu.ng/afnrj

Original Article



Phytochemistry and haematological markers in cadmium-induced toxicity mitigated by date palm (*Phoenix dactylifera* L.) fruit extract in rabbit bucks



Anietie Archibong ANSA¹*^(D), Oyisi Mercy EJIVADE¹^(D) & Odamesan AKPERE²^(D)

¹Department of Animal Science, Faculty of Agriculture, Nnamdi Azikiwe University, Awka, Anambra State, Nigeria ²Department of Animal Science, Faculty of Agriculture, University of Benin, Benin City, Edo State, Nigeria.

DOI: <u>https://www.doi.org/10.5281/zenodo.14020441</u>

Editor: Dr Onyekachi Chukwu, Nnamdi Azikiwe University, NIGERIA

Received: February 2, 2024 Accepted: March 26, 2024 Available online: March 31, 2024

Peer-review: Externally peerreviewed



Copyright: © 2024 Author(s) This is an open access article under the licensed under Creative Commons Attribution 4.0 International License (<u>https://</u> <u>creativecommons.org/licenses/by/</u> <u>4.0/</u>) which permits unrestricted use distribution, and reproduction in any medium, provided the origina author and source are credited.

Conflict of Interest: The authors have no conflicts of interest to declare

Financial Disclosure: The authors declared that this study has received no financial support

ABSTRACT

This study investigated the phytochemical composition of Phoenix dactylifera L. fruit, the effect of cadmium (Cd) on haematological parameters, and the potential of the methanolic extract of Phoenix dactylifera (MEPD) fruit to mitigate Cd-induced toxicity. Forty-five rabbit bucks (24-28 weeks old, 1.41-1.43 kg) were randomized into five groups: control, 3 mg CdCl₂/kg feed, Cd + 300 mg MEPD, Cd + 600 mg MEPD, and Cd + 900 mg MEPD. Cd was administered for 7 days, followed by MEPD treatment every 72 hours for 28 days. The result of phytochemical analysis showed the absence of tannins, phlobatannins, and starch, with a moderate presence of sterols and protein-xanthoproteins. Saponins, terpenoids, cardiac glycosides, alkaloids, phenols, and flavonoids were abundant. Haematological findings revealed no significant differences (P > 0.05) in the Cd-only group compared to the control. MEPD treatment produced a non-significant, dose-dependent increase in packed cell volume, haemoglobin, and red blood cell counts. Significant differences (P < 0.05) were found in mean corpuscular volume and mean corpuscular haemoglobin concentration, though all values remained within the normal range. Platelet counts, significantly (P < 0.05) reduced by Cd (204.30) $\times 10^{3}/\mu$ l), improved significantly with MEPD treatment. This study highlights the phytochemical richness of Phoenix dactylifera and its potential to alleviate Cd-induced haematological toxicity.

K E Y W O R D S : Blood, Date fruit, Heavy metal, Phytotherapy

INTRODUCTION

The escalating environmental pollution presents a significant global concern attributed mainly to the use of harmful chemical substances. Among these, heavy metals, notably implicated in adverse effects even at low concentrations, are of particular concern (Edo *et al.*, 2024. While these metals exist naturally in ecosystems in varying amounts, human activities contribute significantly to their introduction, posing ecological risks

due to their toxicity, persistence, and ability to traverse food chains (Briffa *et al.*, 2020). This contamination, affecting land and water, poses health risks not only to livestock and wildlife but also to various organisms like fish, birds, and humans.

Cadmium (Cd), a prominent heavy metal, has profound implications for public health, being linked to liver enzyme alterations, kidney damage, and long-term impacts on the reproductive, renal, hepatic, and endocrine

*Corresponding author: aa.ansa@unizik.edu.ng; +2348130687781

systems (Fahim et al., 2012; Orisakwe, 2014). In the face of this escalating environmental concern, particularly the adverse impacts of heavy metals like Cd, there is a growing interest in exploring natural remedies which is believed to be a holistic therapy (Karim et al., 2015) when compared to its synthetic counterpart. Phoenix dactylifera, commonly known as the date palm, is a tropical tree known for its rich phytochemicals (Al-Daihan & Bhat, 2012). These inherent bioactive compounds in various parts of the Date palm tree have been reviewed to possess diverse beneficial properties (Al-Okbi, 2022). It was suggested that these properties conferred by Date palm's phytochemicals could potentially serve as an effective therapy against physiological dysfunctions induced by Cd exposure (Zouari et al., 2016; El-Said et al., 2023).

With this perspective, the investigation into the phytochemical screening and haematological parameters becomes imperative in understanding the potential protective effects of the methanolic extract of Phoenix dactylifera fruit against cadmium-induced toxicity. The intricate interplay between the phytochemical components and haematological markers holds the key to unravelling the potential therapeutic benefits of Date palm against the toxic effects induced by cadmium. Therefore, this study aims to assess the phytochemical composition of the methanolic extract of Phoenix dactylifera fruit and evaluate its effect on haematological parameters in rabbit bucks subjected to cadmium exposure.

MATERIALS AND METHODS

Location of study

The experiment was carried out at the Rabbitry Unit of the Teaching and Research Farm in the Department of Animal Science, University of Benin, Benin City, Nigeria. Benin City is located on latitude $6^{\circ}20'17.34''$ N and longitude $5^{\circ}37'32.70''$ E.

Source of Phoenix dactylifera fruits

Dates from the *Phoenix dactylifera* plant were obtained from the Nigerian Institute for Oil Palm Research (NIFOR) in Benin City, Edo State, Nigeria. The acquired fruits were authenticated by a botanist in Department of Plant Biology and Biotechnology, Faculty of Life Sciences, University of Benin, Benin City. The fruits were split, air-dried, finely ground using an electric blender and subsequently stored in an airtight container.

Preparation of fruit extracts

A quantity of 0.5 kg of finely ground *Phoenix dactylifera* fruits was measured and the extraction process was conducted using 99 % methanol in a Soxhlet apparatus. The resulting extract was concentrated with the aid of a



AFNRJ | https://www.doi.org/10.5281/zenodo.14020441 Published by Faculty of Agriculture, Nnamdi Azikiwe University, Nigeria.

rotary vacuum evaporator and freeze dried using a lyophilizer (Wang and Weller, 2006). The freeze-dried extract was then stored in a sealed bottle.

Qualitative phytochemical screening

Qualitative analysis of phytochemicals was conducted on the obtained crude extract, following the established procedures outlined by Harborne (2013). The constituents of interest included tannins, saponins, alkaloids, phenols, flavonoids, sterols, phlobatannins, protein-xathoprotein, cardiac glycosides, terpenoids, and starch.

Acute toxicity evaluation

Acute toxicity of methanolic extract of *Phoenix dactylifera* (MEPD) fruit was studied using the method outlined by Lorke (1983). Twenty-five (20) matured albino rats of both sexes with body weight range of 25 to 30 g were randomized into five groups of four rats per group. Each group received a single dose of NBL extract at 0, 500, 1000, 2000 and 3000 mg/kg body weight, respectively, through oral routes, while the control group received normal saline at 10 ml/kg body weight. The rats were closely observed for signs of toxicity for 72 hours.

Experimental materials and management

A total of forty-five (45) composite rabbit bucks aged 24 - 28 weeks and weighing between 1.41-1.43 kg was used for this study. The rabbits were kept in separate cages under intensive management. A period of two weeks was set aside for quarantine during which the rabbits received Ivomec® injections to control haemoparasites, as well as internal and external parasites. Throughout the study, the rabbits had unrestricted access to both water and commercial grower diet comprising 15 % crude protein and energy content of 2650 kcal/kg.

Experimental design

The treatment protocols consisted of 5 groups: group 1 (control), group 2 (3 mg of CdCl₂/kg feed/day for 7 days), group 3 (CdCl₂/kg feed/day for 7 days + 300 mg/kg body weight of MEPD fruits for 28 days), group 4 (CdCl₂/kg feed/day for 7 days + 600 mg/kg body weight of MEPD fruits for 28 days) and group 5 (CdCl₂/kg feed/day for 7 days + 900 mg/kg body weight of MEPD fruits for 28 days). Each treatment group was replicated three times with three bucks constituting a replicate in a completely randomized design.

Data collection and evaluation

Blood collection: At the end of 35 days research period, 2.5 ml of blood sample was aspirated from each rabbit buck through the ear vein using needle and syringe at 0600 to 0900 hours. The aspirated blood was emptied into sample bottles containing ethylene diamine tetra acetic acid (EDTA) for haematological analysis

Estimation of haematological parameters: Haematological parameters including haemoglobin (Hb) concentration, red blood cell (RBC) count, packed cell volume (PCV), white blood cell (WBC) count, lymphocyte, eosinophils, monocytes, basophils. neutrophils and platelet counts were measured in the whole blood using the automated multi-parameter blood analyser SYSMEX KX21. 50 µl of blood samples was introduced into the equipment and it automatically employed the differences in characteristics possessed by each of the blood components to distinguish and estimate the haematological parameters.

Statistical analysis

The data obtained in this study was subjected to statistical analysis of variance (ANOVA) procedure of GenStat 12th edition at significance level of 5 %. Significant means were separated using Duncan Multiple Range Test (DMRT).

RESULTS AND DISCUSSIONS

The result of qualitative phytochemical screening in this study is summarized in Table 1. The result shows that

saponin, terpenoids, cardiac-glycosides, alkaloids, phenols and flavonoids were highly present in MEPD fruit. The presence of sterols was moderately detected while tannins, phlobatannins and starch were absent.

 Table 1: Phytochemical screening of the methanolic

 extract of *Phoenix dactylifera* fruit

Phyto-constituents	Phoenix dactylifera L.
Tannins	_
Saponins	+ +
Terpenoids	+ +
Sterols	+
Phlobatannins	_
Protein-xathoprotein	+
Cardiac-glycosides	+ +
Alkaloids	+ +
Phenols	+ +
Flavonoids	+ +
Starch	_

++ highly present; + moderately present; - absent.

In the result of haematological study shown in Table 2, there were no significant (P > 0.05) differences between the control and the Cd-only treated group for all the measured parameters.

Table 2:	Haematological profile of	cadmium exposed rabbits administered	Phoenix dactylifera extract
	8 1	1	

Parameters	Control	Cd	Cd + 300 MEPD	Cd + 600 MEPD	Cd + 900 MEPD	SEM
PCV (%)	30.00	30.00	32.67	33.67	35.67	3.53
Hb (g/dl)	8.77	9.10	9.90	10.63	10.83	1.23
RBC (×10 ⁶ /µl)	4.19	4.12	5.11	5.12	5.20	0.62
MCV (fl)	72.23ª	65.77 ^{ab}	64.00 ^b	65.83 ^{ab}	68.73 ^{ab}	2.25
MCH (pg)	20.63	20.17	19.40	20.77	20.83	0.84
MCHC (g/dl)	29.27 ^b	30.70 ^{ab}	30.30 ^{ab}	31.50 ^a	30.30 ^{ab}	0.51
WBC (×10 ³ /µl)	5.13	3.33	5.87	5.13	5.93	0.93
Neutrophils (%)	30.67	35.67	32.00	40.33	44.00	5.76
Lymphocytes (%)	67.00	60.67	64.33	56.00	52.33	5.82
Eosinophils (%)	1.67	2.00	2.00	2.00	2.00	0.39
Monocytes (%)	0.67	1.67	1.67	1.33	1.33	0.80
Basophils (%)	0.00	0.00	0.00	0.33	0.33	0.21
Platelets (×10 ³ /µl)	279.70 ^{ab}	204.30 ^b	361.30 ^a	344.00 ^{ab}	315.30 ^{ab}	44.10

^{a,b} Means bearing different letters of superscript within the same row differ significantly (P < 0.05), PCV- Packed cell volume; Hb- Haemoglobin; RBC- Red blood cell; MCV- Mean corpuscular volume; MCH- Mean corpuscular haemoglobin; MCHC- mean corpuscular haemoglobin concentration; WBC- White blood cell

The administration of MEPD significantly affected (P < 0.05) MCV, MCHC and platelets counts. The MCV value for 300 mg/kg group was significantly (P < 0.05) lower than the control group but similar (P > 0.05) to Cd-only, 600 mg/kg and 900 mg/kg groups. Also, the 600 mg/kg group was significantly (P < 0.05) higher than the control group but similar to Cd-only, 300 mg/kg and 900 mg/kg treated groups. Although the mean values for PCV, Hb,

RBC and WBC counts were not significantly different, these indices presented higher numerical differences in the extract treated groups when compared to both the control and Cd-only treated rabbit bucks. PCV, Hb, RBC, MCV, MCH and neutrophils had a dose dependent effect which progressively increased as the levels of extract administration increases. Only the 600 mg/kg and 900



AFNRJ | https://www.doi.org/10.5281/zenodo.14020441 Published by Faculty of Agriculture, Nnamdi Azikiwe University, Nigeria. mg/kg treated groups recorded a mean value for basophils which was 0.33 % apiece for both treatment groups.

The assessment of the methanolic extract from Phoenix dactylifera fruit revealed the presence of various phytochemical compounds, including tannins, saponins, terpenoids, sterols, phlobatannins, protein-xathoprotein, cardiac glycosides, alkaloids, phenols, flavonoids, and starch. Notably, alkaloids, phenols, flavonoids, saponins, and tannins are important secondary metabolites and have significant role in the medicinal properties of herbs. Previous study on the entire P. dactylifera plant by Vyawahare et al. (2008), indicated the presence of carbohydrates, alkaloids, steroids, flavonoids, vitamins, and tannins. Moreover, investigations into the phytochemical composition of date fruit pulp revealed the abundance of phenolics, sterols, carotenoids, anthocyanins, procyanidins, and flavonoids (Al-Daihan & Bhat, 2012). Additionally, Nadeem et al. (2011) identified oxalate, phytate, and tannin in *P. dactylifera* fruit extract.

Discrepancies observed in the presence of these constituents in *P. dactylifera* across different studies may be attributed to variations in date fruit variety, processing methods, fruit-picking stage, geographical location, and soil conditions. Comparative analyses involving fresh and dried dates have indicated a significant rise in phenolic content during the drying process, likely due to tannin degradation and the maturation of degradative enzymes at elevated temperatures (Al-Farsi *et al.*, 2005). Consequently, the drying process may lead to a reduction in tannin concentration, potentially explaining its absence in the present study.

The assessment of acute oral toxicity in this study revealed that the MEPD fruit caused neither mortality nor observable health concerns in the test rats, even when given at the highest dose of 3000 mg/kg over the 72-hour monitoring period. The rats exhibited typical behaviour, retained their regular feeding patterns, and showed no variation in the consistency of their droppings compared to the control group.

Cd has been reported to induce the formation of metallotioneins and reactive oxygen species (ROS) in the blood, leading to oxidative damage in erythrocytes and various tissues (Kanter *et al.*, 2005). In this study, rabbits that were fed cadmium chloride did not exhibit any significant differences (P > 0.05) in the mean values of erythrocyte counts, haemoglobin (Hb), and packed cell volume (PCV). Moreover, these values aligned with the normal range of $3.48-6.0 \times 10^6$ /mm³ (Chineke *et al.*, 2006; Amata, 2010), 8.9-15.5 g/dl (Poljičak-Milas *et al.*, 2009) and 33-50 % (Fudge, 2000) reported for erythrocyte, Hb and PCV respectively.

Previous research by Hounkpatin *et al.* (2013) on Wistar rats exposed to chronic doses of cadmium chloride at 0.25 and 2.5 mg/kg, given orally for 28 days did not show any significant differences (P > 0.05) in various haematological indices. In contrast, other studies by Dallak (2009) and Onwuka *et al.* (2010) indicated decreased values in erythrocyte counts, Hb, and haematocrit as well as a decreased MCV, MCH and MCHC, suggesting microcytic and hypochromic anaemia in rats orally administered CdCl₂.

In this study, administration of MEPD resulted in a dosedependent non-significant increase (P > 0.05) in PCV, Hb, and RBC. This finding is consistent with the observations of Onuh *et al.* (2012), who found increased levels of RBC, Hb, PCV, reticulocytes, and platelet count in Wistar rats treated with extracts of *Phoenix dactylifera* in a dosedependent manner. Similarly, Wahab *et al.* (2010) reported improved haematological indices in rats with lead acetate-induced haematotoxicity after being administered the ethanolic extract of *Phoenix dactylifera* fruit. This outcome may be related to the phytochemical composition of MEPD, which could have stimulated haematopoietic activity and enhanced bone marrow function.

Erythrocyte indices play a crucial role in the initial categorization of anaemia and serve as an indicator of the bone marrow's ability to generate red blood cells (Brugnara and Mohandas, 2013). The result of erythrocyte indices in this study indicated significant variations (P < 0.05) among the treatment means for mean corpuscular volume (MCV) and mean corpuscular haemoglobin concentration (MCHC). Despite these significant differences, the mean values fell within the normal range of 58.0-79.6 fl and 27-34 g/dl as reported by Poljičak-Milas *et al.* (2009) and Fudge (2000) for MCV and MCHC, respectively. Therefore, the observed differences in MCV and MCHC may not have presented any clinical concerns for the rabbits in the study.

While the Cd-only treated group showed the lowest mean value for white blood cell (WBC) count among all treatment groups, the total and differential WBC counts did not show significant differences (P > 0.05) among the treatment means and were within the established range of $3-12 \times 10^{3}$ /µl, 30-70 %, 30-70 %, 0-3 %, and 0-1 % for WBC, neutrophils, lymphocytes, monocytes, and basophils, respectively, as reported by Hein and Hartmann (2003). Similarly, Wahab et al. (2010) did not observe significant differences in WBC values and its differentials when administering the ethanolic extract of Phoenix dactylifera fruits to rats with lead-induced haematotoxicity.



AFNRJ | <u>https://www.doi.org/10.5281/zenodo.14020441</u> Published by Faculty of Agriculture, Nnamdi Azikiwe University, Nigeria. In this study, although cadmium treatment significantly (P < 0.05) reduced platelet counts, the administration of MEPD significantly (P < 0.05) increased mean platelet counts in the groups treated with the extract. This supports the conclusions of Ramadhas *et al.* (2014), suggesting that *Phoenix dactylifera* extract may be effective in correcting bleeding disorders associated with thrombocytopenia.

CONCLUSION AND RECOMMENDATION

In conclusion, the study indicates that the methanolic extract from *Phoenix dactylifera* fruit contains various phytochemicals and may have a protective effect on haematological parameters in rabbits exposed to cadmium chloride, potentially due to its haemopoietic activity. It is recommended that *Phoenix dactylifera* fruit extract, particularly at doses of 600 mg/kg and 900 mg/kg, be further investigated for its potential to mitigate cadmium-induced haematotoxicity and enhance haematopoietic function.

Acknowledgments

The authors express their appreciation to the Department of Chemistry, Faculty of Physical Sciences, University of Benin, Benin City, Nigeria, for granting access to their facilities during this research. Additionally, they acknowledge the Department of Plant Biology and Biotechnology, University of Benin, Benin City, Nigeria for the assistance in identifying the fruit used in this study.

Authors Contribution

Research Design: AAA; Data analysis/interpretation: AAA; Data Collection: OA; Literature Search: OME; Writing/proofreading: AAA & OME. All authors read and approved the final manuscript.

Ethical Committee Approval

This experiment was approved and conducted according to the provisions of the Ethical Committee on the use of animals for biomedical research at the University of Benin, Benin City, Nigeria (Approval No: PG/AGR1312015).

REFERENCES

- Al-Daihan, S. & Bhat, R. S. (2012). Antibacterial activities of extracts of leaf, fruit, seed and bark of *Phoenix dactylifera. African Journal of Biotechnology*. 11(42): 10021-10025. <u>https://doi.org/10.5897/ajb11.4309</u>
- Al-Farsi, M., Alasalvar, C., Morris, A., Baron, M. & Shahidi, F. (2005). Comparison of antioxidant activity, anthocyanins, carotenoids, and phenolics of three native fresh and sun-dried date (Phoenix dactylifera L.) varieties grown in Oman. Journal of

Agricultural and Food Chemistry. 53(19):7592–7599. https://doi.org/10.1021/jf050579q

- Al-Okbi, S.Y. (2022). Date Palm as Source of Nutraceuticals for Health Promotion: a Review. Current Nutrition Reports. 11: 574–591. <u>https://doi.org/10.1007/s13668-022-00437-w</u>
- Amata, A. (2010). The effect of feeding *Gliricidia* leaf meal (GLM) on the haematological, serological and carcass characteristics of weaned rabbits in the tropics. *Agriculture and Biology Journal of North America*. 1(5): 1057-1060. https://doi.org/10.5251/abjna.2010.1.5.1057.1060
- Briffa, J., Sinagra, E. & Blundell, R. (2020). Heavy metal pollution in the environment and their toxicological effects on humans. *Heliyon*, 6(9): e04691. <u>https://doi.org/10.1016/j.heliyon.2020.e04691</u>
- Brugnara, C. & Mohandas, N. (2013). Red cell indices in classification and treatment of anemias: from M.M. Wintrobes's original 1934 classification to the third millennium. *Current Opinion in Hematology*. 20(3): 222-230. https://doi.org/10.1097/moh.0b013e32835f5933
- Chineke, C.A., Ologun, A.G. & Ikeobi, C.O. (2006). Haematological parameters in rabbit breads and crosses in humid tropics. *Pakistan Journal of Biological Sciences*. 9(11): 2102-2106. https://doi.org/10.3923/pjbs.2006.2102.2106
- Dallak, M. (2009). Camel's milk protects against cadmium chloride induced hypocromic microcytic anaemia and oxidative stress in red blood cells of white albino rats. *American Journal of Pharmacology and Toxicology*. 4(4): 136-143. <u>https://doi.org/10.3844/ajptsp.2009.136.143</u>
- Edo, G.I., Samuel, P.O., Oloni, G.O., Ezekiel, G.O., Ikpekoro, V.O., Obasohan, P., Ongulu, J., Otunuya, C.F., Opiti, A.R., Ajakaye, R.S., Essaghah, A.E.A. & Agbo, J. J. (2024). Environmental persistence, bioaccumulation, and ecotoxicology of heavy metals. *Chemistry and Ecology*, 40(3), 322–349. https://doi.org/10.1080/02757540.2024.2306839
- El-Said, K., Amoush, W. & Mohamed, A. (2023). Effect of Phoenix dactylifera seeds extract on cadmiuminduced hepatotoxicity in male mice. *Journal of Bioscience and Applied Research*, 9(1), 53-61. <u>https://doi.org/10.21608/jbaar.2023.285435</u>
- Fahim, M. A., Nemmar, A., Dhanasekaran, S., Singh, S., Shafiullah, M., Yasin, J., Zia, S. & Hasan, M. Y. (2012). Acute cadmium exposure causes systemic and thromboembolic events in mice. *Physiological Research*.61(1):73-

80. https://doi.org/10.33549/physiolres.932238

- Fudge, C.S (2000) Laboratory Medicine: Avian and Exotic pets. WB Sanders, Philadelphia, WB Saunders. pp 9–18.
- Harborne, J. B. (2013). Phytochemical methods : a guide to modern techniques of plant analysis. 18th ed. New Delhi (India) : Springer. pp 135-203.



AFNRJ | <u>https://www.doi.org/10.5281/zenodo.14020441</u> Published by Faculty of Agriculture, Nnamdi Azikiwe University, Nigeria.

- Hein, J. & Hartmann, K. (2003). Labordiagnostiche Referez-bereiche bei Kaninchen. *Tieraerzti prax* 31(5): 321-328. <u>https://doi.org/10.1055/s-0037-1622371</u>
- Hounkpatin, A. S. Y., Edorh, P. A., Guédénon, P., Alimba, C. G., Ogunkanmi, A., Dougnon, T. V., Boni, G., Aissi, K. A., Montcho, S., Loko, F., Ouazzani, N., Mandi, L., Boko, M. & Creppy, E. E. (2013). Haematological evaluation of Wistar rats exposed to chronic doses of cadmium, mercury and combined cadmium and mercury. *African Journal* of *Biotechnology*. 12(23): 3731-3737. https://doi.org/10.5897/AJB12.2669
- Kanter, M., Coskun O. & Gurel, A (2005). Effect of black cumin (*Nigella sativa*) on cadmium-induced oxidative stress in the blood of rats. *Biological Trace Element Research*. 107(3): 277-287. <u>https://doi.org/10.1385/bter:107:3:277</u>
- Karimi, A., Majlesi, M., & Rafieian-Kopaei, M. (2015). Herbal versus synthetic drugs; beliefs and facts. *Journal of nephropharmacology*, 4(1), 27–30.
- Lorke, D. (1983). A new approach to practical acute toxicity testing. Archives of toxicology, 54(4): 275-287. <u>https://doi.org/10.1007/bf01234480</u>
- Nadeem, M., Rehman, S. U., Anjum, F. M., Zahoor, T., Saeed, F. & Ahmad, A. (2011). Anti-nutritional Factors in Some Date Palm (*Phoenix dactyliferaL.*) Varieties Grown in *Pakistan. Internet Journal of Food Safety*.13: 386-390.
- Onuh, S. N., Ukaejiofo, E. O., Achukwu, P. U., Ufelle, S. A., Okwuosa, C. N. & Chukwuka, C. J. (2012). Haemopoietic activity and effect of crude fruit extract of *Phoenix dactylifera* on peripheral blood parameters. *International Journal of Biological & Medical Research.* 3(2): 1720-1723.
- Onwuka, F.C., Erhabor, O., Eteng, M.U. & Umoh, I.B (2010). Ameliorative effect of cabbage extract on cadmium induced changes on hematology and biochemical parameters of albino rats. *Journal of Toxicology and Environmental Health Sciences*. 2(2):11-16.

- Orisakwe, O. E. (2014). Lead and Cadmium in public health in Nigeria: Physicians Neglect and Pitfall in Patient Management. North American Journal of Medicine and Science. 6(2): 61–70. https://doi.org/10.4103/1947-2714.127740
- Poljičak-Milas, N., Kardum-Skelin, I., Vuđan, M., Silvija Marenjak, T., Ballarin-Perharić, A. & Milas, Z. (2009). Blood cell count analyses and erythrocyte morphometry in New Zealand white rabbits. *Veterinarski arhiv*, 79(6): 561-571. https://hrcak.srce.hr/53081
- Ramadhas, M., Palanisamy, K., Sudhagar, M. & Mani, V. M. (2014). Ameliorating effect of Phoenix dactylifera on lambda cyhalothrin induced biochemical, hematological and hepatopathological alterations in male wistar rats. Biomedicine & Aging Pathology. 4(3): 273–279. https://doi.org/10.1016/j.biomag.2014.04.002
- Vyawahare, N., Pujari, R., Khsirsagar, A., Ingawale, D., Patil, M., & Kagathara, V. (2008). *Phoenix dactylifera*: An update of its indegenous uses, phytochemistry and pharmacology. *The Internet Journal of Pharmacology*. 7(1): 1-11. https://doi.org/10.5580/164b
- Wahab, A. A., Mabrouk, M. A. A., Joro, J. M., Oluwatobi, S. E., Bauchi, Z. M. & John, A. A. (2010). Ethanolic extract of *Phoenix dactylifera* L. prevents lead induced hematotoxicity in rats. *Continental Journal* of Biomedical Sciences. 4: 10-15.
- Wang, L. & Weller, C. (2006) Recent advances in extraction of nutraceuticals from plants. *Trends in Food Science and Technology*, 17(6), 300 – 312 <u>http://dx.doi.org/10.1016/j.tifs.2005.12.004</u>
- Zouari, M., Elloumi, N., Ahmed, C. B., Delmail, D., Rouina, B. B., Abdallah, F. B. & Labrousse, P. (2016). Exogenous proline enhances growth, mineral uptake, antioxidant defense, and reduces cadmium-induced oxidative damage in young date palm (*Phoenix dactylifera* L.). Ecological Engineering, 86, 202-209. https://doi.org/10.4314/jasem.v26i5.12



AFNRJ | https://www.doi.org/10.5281/zenodo.14020441 Published by Faculty of Agriculture, Nnamdi Azikiwe University, Nigeria.