

REVIEW OF LABORATORY BIOMARKERS OF INFLAMMATORY BOWEL DISEASE

Authors:

Udegbune, M. M

Author Affiliation:

Department of Biomedical Science and Physiology, University of Wolverhampton, U.K

Corresponding Author:

Michael M Udegbune

Department of Biomedical Science and Physiology,

University of Wolverhampton,

Wolverhampton, U.K.

E-mail: mmadukaudegbune@yahoo.com

XX

ABSTRACT

Inflammatory bowel disease (IBD), a cluster of chronic, immune-related and long-term disorders that cause fever, abdominal pain/cramping, fatigue, severe and recurrent diarrhoea, rectal bleeding and weight loss, and characterised by inflammation of gastrointestinal tract (GIT), is categorised as crohn's disease (CD), ulcerative colitis (UC) or IBD unclassified (IBDU). Assessing the diagnosis, severity and monitoring of IBD is largely based on the combined effects of clinical presentations, endoscopy, radiology, histology and laboratory biomarkers. Some laboratory biomarkers of IBD such as C-reactive protein (CRP), erythrocyte sedimentation rate (ESR), platelets etc are associated with systemic and gastrointestinal inflammation or disease activity, i.e., active or quiescent IBD. Others are linked to genetic predisposition, e.g., autophagy genes, nucleotide-binding oligomerization domain-containing protein 2 (NOD2), interleukin-23 receptor (IL23R) etc, correlated with the neoplastic transformation, e.g., M2-pyruvate kinase, miRNAs, mucosal chitinase-3-like protein 1 (CH13L1) etc, and coupled to drug metabolism in relation to therapeutic intervention e.g., Thiopurine Methyltransferase (TPMT) and 6-Thioguanine Nucleotide (6TGN). There are also those such as anti-neutrophil cytoplasmic antibodies (ANCA), and anti-saccharomyces cerevisiae antibodies (ASCA) that are related to the type of IBD, i.e., either CD, UC or IBDU. This review summarises the characterisation of laboratory biomarkers of IBD including a narrative on their future perspectives from the standpoint of diagnosis, prognosis, monitoring the disease course and a practical algorithm for the practical application of these biomarkers in the assessment of IBD.

Keywords: Biomarker, Inflammatory Bowel Disease, Crohn's Disease, Ulcerative Colitis, S100 Proteins, Calprotectin, S100A12, C-reactive Protein.

INTRODUCTION

Inflammatory bowel disease (IBD), a cluster of chronic, immune-related and long-term disorders that cause fever, abdominal pain/cramping, fatigue, severe and recurrent diarrhoea, rectal bleeding and weight loss, and characterised by inflammation of the gastrointestinal tract (GIT), is categorised as crohn's disease (CD), ulcerative colitis (UC) or IBD unclassified (IBDU).¹⁻³

Epidemiologically, IBD has witnessed considerable changes with consequent annual variations in the incidence by geographical region and increase in the disease burden globally. With worldwide prevalence at ≈ 5 million, the incidence per 100,000 population in Europe stood at ≈ 10.5 – 46.1% , Middle East at ≈ 23.7 – 39.8% , North America at ≈ 7.3 – 0.2% , Oceania at ≈ 0.21 – 3.67% and Asia at ≈ 1.37 – 1.5% , there appears to be an increasing trend with age in the disease burden in adults (exceeding 1% in those >70 years) relative to children and adolescents, and in female (≈ 3.9 million) compared to male (≈ 3.0 million) population.¹⁻³

The advent of more convenient, available and affordable laboratory biomarkers for the diagnosis, prognosis and treatment of IBD is expected to totally change the way that patients presenting with the disease are treated¹. Although the recent scientific and technological advances in the development, optimisation and analytical validation of immunoassay kits have somewhat eased the practical issues associated with the application of laboratory biomarkers in the assessment of IBD, there still exist some uncertainties over the gains recorded with the advances in genomic, proteomic and metabolics arrays in relation to clinical laboratory testing¹.

Recent advances in immunoassay technologies have necessitated the categorisation of these laboratory biomarkers of IBD into those biomarkers that are associated with systemic and gastrointestinal inflammation including disease activity, i.e., active or quiescent IBD; those biomarkers that are linked to genetic predisposition of IBD; those biomarkers that are correlated with the neoplastic transformation of IBD; those biomarkers that are coupled to drug metabolism in relation to therapeutic intervention of IBD; and those biomarkers that are related to type of IBD, i.e., either CD, UC or IBDU.¹⁻³

More information may be needed on the many aspects of laboratory biomarkers to diagnose IBD, differentiate between disease states (CD vs. UC), evaluate disease activity (active vs. quiescent), confirm disease sites or locations as in small intestine: ileal and upper (A1) vs. colonic and ileo-colonic (A2); large intestine: proctitis (E1) vs. distal colitis (E2) vs. pancolitis (E3), predict the disease course and relapse, and monitor follow-up and response to treatment.¹

The diagnosis, severity and monitoring of IBD are usually based on a combination of clinical presentations, endoscopy, radiology, laboratory biomarkers and, if appropriate histology. Laboratory biomarkers should be non-invasive, cheap, simple, objective, rapid, easy to perform and reproducible (*table 1*). An ideal biomarker for IBD that would combine all these characteristics is not available.^{1,2} Serum biomarkers of systemic inflammation are shown in *table 2* and faecal biomarkers of gastrointestinal inflammation are shown in *table 3*.

Faecal biomarkers may be regarded as the 'gold standards' in intestinal inflammation. They are however, limited by certain factors including

lack of specificity for IBD; lack of validated, optimal and varying assay cut-offs make it difficult to characterize active inflammatory disease, distinguish IBD from irritable bowel syndrome (IBS), predict clinical remission and mucosal healing, and assess response to treatment; levels of calprotectin in stool samples are dependent on diverse physiological considerations such as age and clinical comorbidities; significant overlap in faecal calprotectin levels (50–150 µg/g) in both IBD and IBS patients presents an ambiguous situation regarding the decision to refer a patient to endoscopy or not, and intra-individual and inter-individual variability, spot variability in the same sample and reluctance of some patients to provide stool samples.¹⁻³ These issues may be overcome by using serum samples for the measurement of calprotectin and S100A12.¹ This offers the prospect that serum calprotectin and serum S100A12 could replace or supplement faecal calprotectin and faecal S100A12 in the identification and assessment of IBD.

Whilst it is important for laboratory biomarkers of IBD to be non-invasive, cheap, simple, objective, rapid, easy to perform and reproducible, the 'real-world' experience with these biomarkers is increasing together with the increasing number of patients with IBD for which the usefulness of these laboratory biomarkers to assess the disease have become necessary.¹⁻³ The 'ideal biomarker' that would integrate all the preceding characteristics for evaluating the analytical performance and diagnostic accuracy of these laboratory biomarkers in IBD is not available.¹ Therefore, the search for the availability of a single laboratory biomarker that would demonstrate the attributes of an 'ideal biomarker' for use in assessing IBD continues.

This objectives of this review were to present a summary of the characterisation of laboratory biomarkers of IBD including a narrative on their future perspectives from the standpoint of diagnosis, prognosis and monitoring the disease course. A practical algorithm for the practical application of these biomarkers, both current and potential future biomarkers, in the assessment of IBD will also merit discussion. A conclusion will be added.

A. Laboratory biomarkers with widespread application

1. Serum C-reactive protein and cytokines

C-Reactive Protein (CRP), a pentameric protein synthesized predominantly in the liver but also in vascular walls and adipose tissue, is a well-established biomarker of acute inflammation. During acute phase response, hepatocytes rapidly increase production of CRP under the influence of interleukin-6 (IL-6), tumour necrosis factor-alpha (TNF-α), and interleukin-1β (IL-1β). Serum CRP, therefore, rises rapidly up to 100 to 1000-fold, peaking in two days and then decreases rapidly with the resolution of the acute insult as it has a half-life of 19 hours. Functionally, CRP is recruited into the complement activation process when it binds to organisms or particles that contain phosphocholine which enables it to opsonise infectious agents and damaged cells.⁴⁻⁶ In healthy individuals, CRP circulates in low concentration (about 1 mg/L). CRP concentrations around 10 to 40 mg/L may be seen in chronic inflammation.^{7,8}

CRP is generally increased in IBD, but appears to respond differently to inflammation in UC and CD. A rise in serum CRP level correlates well with the disease activity in CD. CRP, however, may not be raised or mildly raised in

the presence of increased disease activity in UC even though increased levels of IL-6, IL-1 β or TNF- α are observed.⁹ Other inflammatory markers such as β_2 -microglobulin correlate better with histology scores in UC.^{10,11} The main reason given for this differential CRP response in IBD is that inflammation is limited to the mucosa in UC and, less likely to provoke a

systemic response to inflammation when compared to transmural inflammation in CD. Other possible reasons include increased IL-6 levels in CD compared to UC¹² and CRP gene polymorphism in UC and CD.¹³⁻¹⁵ CRP levels in IBD patients, however, are not associated with CRP gene polymorphism.¹⁶

Table 1 – Characteristics of an ideal biomarker for use in the assessment of IBD¹

Performance	Criteria Characteristics
Simple	Disease specific: Identify individuals with IBD; Able to differentiate IBD from non-IBD cases; Able to predict the remission or relapse; Monitor the effect of treatment; Prognostic value in assessing morbidity/mortality
Ease of performance	Ability to objectively measure disease activity without ambiguity
Invasiveness	Not invasive or minimally invasive
Cost	Affordable
Rapid	Quick turnaround time
Reproducibility	Assay results not showing discrepancies between individuals and clinical laboratories.

Table 2 – Serum acute phase proteins and their responses to IBD and Other inflammatory processes¹. * α_2 -macroglobulin shows a different response in animals in comparison to humans, both in

Acute Phase Protein	Increased (\uparrow)	Decreased (\downarrow)
Proteinase Inhibitors	α_1 -Antitrypsin, α_1 -Antichymotrypsin, α_2 -Macroglobulin*	
Coagulation and Fibrinolytic Proteins	Fibrinogen, Prothrombin, Factor VIII, Plasminogen, Tissue Plasminogen Activator Antithrombin	Factor XII
Complement System	C1s, C2, C3, C4, C5, C1 Inhibitor, C9	Albumin, Transferrin
Transport Proteins	Haptoglobin, Haemopexin, Caeruloplasmin	Insulin-like Growth Factor, α - Fetoprotein, Cholinesterase
Other Acute Phase Proteins	C-reactive protein, Serum Amyloid A, Ferritin, Fibronectin, Orosomuroid (α_1 - Acid Glycoprotein).	

positive and negative acute phase protein respectively.

Table 3 – Faecal biomarkers in clinical use for the diagnosis and differential diagnosis of IBD including its discrimination from IBS³.

Faecal Biomarker	Major Source or Origin
Calprotectin (S100A8/S100A9)	Neutrophils, Monocytes and Epithelial Cells.
Calgranulin C or EN-RAGE (S100A12)	Neutrophils
Lactoferrin	Mucosal Epithelial Cells and Neutrophils
M2-Pyruvate Kinase (M2PK)	Expressed by rapidly dividing cells
Neopterin	Activated Macrophages
Metalloproteinases (MMPs)	Different Cell types including Activated Neutrophils
Myeloperoxidases (MPOs)	Activated Neutrophils
Polymorphonuclear Elastase (PMN)	Activated Neutrophils

CRP is generally increased in IBD, but appears to respond differently to inflammation in UC and CD. A rise in serum CRP level correlates well with the disease activity in CD. CRP, however, may not be raised or mildly raised in the presence of increased disease activity in UC even though increased levels of IL-6, IL-1 β or TNF- α are observed.⁹ Other inflammatory markers such as β_2 -microglobulin correlate better with histology scores in UC.^{10,11} The main reason given for this differential CRP response in IBD is that inflammation is limited to the mucosa in UC and, less likely to provoke a systemic response to inflammation when compared to transmural inflammation in CD. Other possible reasons include increased IL-6 levels in CD compared to UC¹² and CRP gene polymorphism in UC and CD.¹³⁻¹⁵ CRP levels in IBD patients, however, are not associated with CRP gene polymorphism.¹⁶

The utility of CRP as a serum biomarker in the assessment of IBD is still unclear. CRP is useful in assessing IBD patients with the active disease. However, there is the disadvantage that the inclusion of CRP in IBD assessment carries the increased risk of being restricted to IBD patients with high concentration of the protein.¹

Although the application of CRP may offer the promise of an increased prospect of response to treatment with a reduced rate for placebo considerations, it is however, important to emphasise the fact that the United States Food and Drug Administration (FDA) placed restrictions on using CRP in IBD assessment.¹ CRP presents a fair evaluation of the follow-up effect of therapeutic intervention in IBD patients. The ambiguity of CRP cut-off values in assessing IBD makes the utility of CRP very problematic.¹

2. Circulating haemopoietic biomarkers: erythrocyte sedimentation rate (ESR), platelet count, mean platelet volume (MCV) and red cell distribution width (RDW)

ESR is an indirect measure of systemic inflammation. ESR measures the rate of migration of erythrocytes through the plasma, and is increased in the presence of increased proteins including acute phase proteins. ESR, compared to CRP, is slow to peak in response to inflammation and slow to decline after resolution of inflammation and is therefore of limited value in IBD assessment.^{1,17}

Inflammation processes, including IBD, cause an increase in platelets and changes in their morphology. Platelet count increases in patients with IBD, particularly in UC when reticulated platelets are taken into account.^{18,19} It, however, is not a useful biomarker of IBD, given the wide range of a normal platelet count.^{1,20}

MPV has been reported as decreased in active IBD and has a negative correlation with CRP, while other studies have found no correlation between a fall in MPV and the disease activity.^{21,22} Likewise, leucocytes lack specificity as a biomarker for IBD,²³ and are influenced by treatments in IBD with drugs such as glucocorticoids or azathioprine (increased) and 6-mercaptopurine (decreased).¹

RDW is a measure of size variability and heterogeneity of erythrocytes in peripheral blood. RDW increases in active IBD, and particularly in CD compared to UC. RDW at a diagnostic cut-off value of 13.8% in non-anaemic UC patients has a sensitivity and specificity of 76% and 86% respectively, and in non-anaemic CD patients at a diagnostic cut-off value of 14.1% has a sensitivity and specificity of 82% and 83% respectively.^{23,24}

3. Other serum acute phase proteins

Other acute phase reactants (Table 1.3) such as fibrinogen, sialic acid, α_1 -acid glycoprotein (orosomucoid), α_1 -antitrypsin, α_2 -globulin, β_2 -microglobulin, serum amyloid A (SAA) and albumin as biomarkers of IBD have not been widely studied because of their apparent inferiority to CRP. α_1 -acid glycoprotein, shows good correlation with IBD disease activity but its half-life of 5 days makes it unsuitable as IBD biomarker.^{1,17}

4. Faecal lactoferrin

Lactoferrin is an iron-binding protein found in neutrophil granulocytes. In acute intestinal inflammation, increased mucosal infiltration of neutrophils and subsequent secretion of lactoferrin into the intestinal tract results in increased faecal lactoferrin.^{25,26} Lactoferrin has anti-bacterial activity by limiting availability of iron, and causes direct damage to bacterial cell membrane. Faecal lactoferrin is resistant to degradation and proteolysis, although less so than calprotectin, making it a useful biomarker of intestinal inflammation. The significant proportion of faecal lactoferrin in stool is, therefore, a reflection of intestinal inflammation.²⁷⁻²⁹

Lactoferrin is stable in stool for up to 5 days, does not deteriorate with repeated freezing and thawing, and can be quantitatively measured by the enzyme-linked immunosorbent assay (ELISA) technique. It is a non-specific intestinal biomarker being raised not only in active IBD but in other inflammatory intestinal disorders including infective diarrhoea, colon cancer and non-steroidal anti-inflammatory drugs (NSAIDs) enteropathy. Faecal lactoferrin correlates well with the clinical, endoscopic and histological grading of IBD disease activity and therefore, may distinguish between active and inactive IBD, and between IBD patients and healthy controls.^{26,30-36} For predicting relapse, faecal lactoferrin has 46% sensitivity and 61% specificity for UC, and 77% sensitivity and 68% specificity for CD.³⁷ Comparable diagnostic accuracy of lactoferrin and calprotectin exist in patients with IBD and in patients with irritable bowel syndrome (IBS) and healthy controls³⁸. Calprotectin and lactoferrin had similar sensitivity (78% vs. 80%), specificity (83% vs. 85%), overall diagnostic accuracy of 80% vs. 81% and are both significantly elevated in children with IBD.^{38,39}

5. *Faecal neopterin*

Neopterin is a metabolite of biopterin released by activated macrophages and monocytes. Faecal neopterin correlates well with disease activity in IBD particularly with the severity of mucosal lesions. Faecal neopterin shows better correlation with endoscopic scores in UC ($r = 0.72$; $p < 0.0001$) than in CD ($r = 0.47$; $p < 0.0001$). Faecal neopterin is significantly higher in clinically and endoscopically active IBD compared to inactive IBD. The diagnostic accuracy of faecal neopterin to predict endoscopic activity in IBD compares favourably with faecal calprotectin in CD (74%) and in UC (87%) and, like calprotectin and lactoferrin, release of neopterin is non-specific as it could be triggered in response to disease conditions of viral infection and cell-mediated immune response during the early phase of inflammation.^{40,41}

6. *Faecal metalloproteinases (MMPs)*

MMPs are zinc-dependent endopeptidases secreted by various cell types. MMP-9 is the MMP released in highest concentrations by activated neutrophils during intestinal inflammation such as in IBD. While MMP-1, MMP-2 and MMP-3 are detected in significantly higher amount in UC, only MMP-9 levels are significantly higher in active UC than in IBS or healthy controls thereby making it a reliable biomarker in distinguishing between UC and IBS based on 85% sensitivity and 100% specificity.⁴² Faecal MMP-9 correlates well with other faecal biomarkers including calprotectin, lactoferrin and neopterin in endoscopically assessed disease activity in IBD and also demonstrates comparable diagnostic accuracy with faecal calprotectin.⁴³

7. *Faecal myeloperoxidases (MPOs)*

MPOs are lysosomal proteins released by activated neutrophils in response to inflammation. The diagnostic accuracy of faecal MPOs to assess endoscopic disease activity in IBD is inferior to calprotectin and polymorphonuclear (PMN) elastase.⁴⁴

8. *Faecal polymorphonuclear (PMN) elastase*

PMN elastase is released by activated neutrophils during inflammation, and is stable for up to 4 days in stool at ambient temperature. Faecal PMN elastase may be a useful biomarker in IBD as it is significantly higher in patients with active IBD compared to inactive IBD. This is reflected in its diagnostic accuracy of 84% sensitivity and 87% specificity for IBD that increases to 96% sensitivity and 100% specificity when combined with calprotectin and lactoferrin.^{35,45}

9. *Faecal M2-pyruvate kinase (M2PK)*

Faecal levels of M2PK, a multi-functional protein found in undifferentiated and proliferating cells, are associated with active IBD, and correlate well with calprotectin in distinguishing IBD from IBS. M2PK is a superior biomarker to calprotectin, lactoferrin and S100A12 in predicting steroid refractoriness in severe paediatric UC. It also holds promise as a potential screening biomarker for colorectal cancer (CRC) in UC.⁴⁶⁻⁴⁹

10. *The S100 Proteins: Faecal and serum S100A12*

The family of low molecular weight S100 proteins or calgranulins (*figure 1*) have several functions including roles in cellular inflammation, proliferation, differentiation, apoptosis, signal transduction, calcium homeostasis and energy metabolism.⁵⁰⁻⁵³ In chronic inflammation, the S100 protein family

are actively expressed in activated granulocytes.⁵⁴⁻⁵⁶

The cytokine-like extracellular functions of S100 proteins such as chemotactic activities related to inflammation and the acute phase response are exhibited mainly by S100A8 (calgranulin A), S100A9 (calgranulin B) and S100A12 (calgranulin C), and these are commonly referred to as the calgranulins or myeloid-related proteins (MRP).⁵⁷⁻⁵⁹ The name 'S100 proteins' was derived from their ability to be 100% soluble in a saturated solution of ammonium sulphate at neutral pH. These proteins bind calcium, and are characterized by two calcium-binding motifs called the elongation factor (EF) hand. There are over 600 members of EF-hand super family.⁶⁰⁻⁶³ The S100 proteins are expressed in myeloid cells including neutrophils, monocytes, and dendritic cells.⁶⁴

S100A12 (*figure 2*) is a 10.4 kilodalton (*kDa*) molecular weight, 91-amino acids protein. It is a calcium-binding, proinflammatory protein predominantly expressed and secreted by neutrophil granulocytes, and represents 5% of total cytosolic protein.^{65,66} S100A12 is a ligand for the receptor for advanced glycation end products (RAGE) and therefore, also named as extracellular newly identified receptor for advanced glycation end products binding protein (EN-RAGE). Other alternative names for S100A12 are calgranulin C, migration inhibition factor-related protein 6 (MRP-6) and calcium-binding protein in amniotic fluid (CAAF-1).^{57,67,68} The binding of S100A12 to RAGE modulates these extracellular functions in disease. In inflammatory disease, serum S100 correlates with disease activity parameters and in these conditions S100 proteins may be superior to conventional laboratory biomarkers of inflammation including CRP and ESR.⁶⁹⁻⁷¹

Serum S100A12 have been reported in increased concentrations in various biological materials including synovial fluid, synovial tissue and serum of patients with inflammatory arthritis.⁷²⁻⁷⁵ Serum S100A12 is also raised in neurodegenerative diseases, diabetes mellitus, rheumatoid arthritis, osteoarthritis, cancerogenesis, familial mediterranean fever, idiopathic pulmonary fibrosis, cardiovascular diseases and atherosclerosis.⁷⁶⁻⁸² S100A12 levels were also elevated in patients with peripheral radiographic features ($p = 0.036$), but did not correlate with clinical variables of disease activity in psoriatic arthritis.⁸³ In haemodialysis patients, levels of S100A12 are linked to cardiovascular disease.^{84,85} Elevated concentrations of S100A12 and its receptors are found in pulmonary tissue and broncho-alveolar lavage fluid in acute lung injury.^{86,87} Both S100A8 and S100A9 hetero-complexes are actively expressed preceding prostate tumour genesis and subsequent development, progression and enlargement of prostate carcinomas. Alone, serum S100A9 is increased significantly in prostate cancer patients compared to healthy controls or patients with benign prostatic hyperplasia (BPH); thereby underlining its role as a useful biomarker in differentiating prostate carcinoma and BPH.⁸⁸

Faecal S100A12 have been shown to be a novel non-invasive biomarker of IBD in paediatric populations.^{89,90} S100A12 levels in stool can be used as an indicator of disease activity in chronic IBD and to gauge the degree of gastrointestinal tract inflammation. As a biomarker of neutrophil activation, faecal S100A12 could play a significant role as a non-invasive biomarker of intestinal inflammation.




Figure 1 – Dimer structure of S100 protein. S100 proteins are small proteins with a molecular weight of 10 – 12 kDa. Each S100 protein consists of two EF-hand helix-loop-helix structural motifs, which are arranged in a back-to-back manner and linked with a flexible hinge.⁹⁵




Figure 2 – Structure of S100A12. A three-dimensional model of the crystal structure of the S100A12 protein in a hexameric form and its proposed role in receptor signalling. The stereo view of the Ca^{2+} and Cu^{2+} -S100A12 dimer in ribbon representation, with individual subunits shown in red and blue. The chain topology, subunit arrangement, and juxtaposition of metal-binding sites are typical of metal-bound S100 proteins.⁹¹⁻⁹⁴

11. Faecal and serum calprotectin

Calprotectin (figure 3), is a 36 kilodalton (kDa) calcium- and zinc-binding protein composed of two heavy 14 kDa S100A9 (MRP14) and one light 8 kDa S100A8 (MRP8) subunits which are members of the EF-hand motif containing S100 family of proteins. Like all S100 proteins, the genes that code for calprotectin are located within the gene cluster on chromosome 1q21 region. Calprotectin is expressed predominantly in neutrophils and monocytes – in which it constitutes up to 60% of the cytosolic protein. It

is also expressed in macrophages, keratinocytes, epithelial cells and endothelial cells.^{50,96-100}

Calprotectin was first characterised in 1980 as the Leucocyte protein candidate 1 (L1), to reflect *in-vivo* granulocyte turnover.¹⁰¹⁻¹⁰⁴ L1 was renamed calprotectin in recognition of its antimicrobial activity¹⁰⁵, calcium-binding roles and subsequent involvement in intracellular signal transduction and regulatory functions in acute phase response and inflammatory processes^{96,106,107}. Neither S100A8 nor S100A9 subunits in isolation have anti-microbial

characteristics. It is possible that the high affinity of calprotectin subunits, S100A8 and S100A9, for zinc-binding site could explain the reduction of zinc concentration sufficiently to allow calprotectin to inhibit microbial growth.^{106,108-111}

Alternative names for calprotectin include: MRP-8 and MRP-14 (MRP8/MRP14), p8/14, p34 and S100A8/S100A9. Calprotectin is composed of two heterocomplexes: S100A8 (also known as MRP-8, Calgranulin A and CP-10 in mouse) and S100A9 (also known as MRP-14 and Calgranulin B).

Calprotectin is found in plasma, saliva, cerebrospinal fluid, urine and faeces. Circulating calprotectin is commonly raised in inflammatory diseases including acute coronary syndromes, cystic fibrosis, multiple sclerosis, human immunodeficiency virus, rheumatoid arthritis, reactive arthritis, juvenile chronic arthritis, juvenile idiopathic arthritis, psoriatic arthritis, polymyalgia rheumatica, systemic lupus erythematosus and acute rejection in kidney allograft transplantation^{103,112-130}

Calprotectin is found in plasma, saliva, cerebrospinal fluid, urine and faeces. Circulating calprotectin is commonly raised in inflammatory diseases including acute coronary syndromes, cystic fibrosis, multiple sclerosis, human immunodeficiency virus, rheumatoid arthritis, reactive arthritis, juvenile chronic arthritis, juvenile idiopathic arthritis, psoriatic arthritis, polymyalgia rheumatica, systemic lupus erythematosus and acute rejection in kidney allograft transplantation^{103,112-130}

Calprotectin expression correlates with microglial activation in cerebral malaria; and serum levels are prognostic biomarkers in recurrent infection and survival in alcoholic liver disease.¹³¹⁻¹³³ Increased calprotectin concentrations in faecal samples of patients with

IBD that reflect granulocyte migration through the inflamed intestinal wall has been reported.¹³⁴⁻¹³⁶ Measurement of faecal calprotectin now enjoys wide application in the diagnosis and monitoring of IBD, and an increased concentration indicates organic intestinal disorders. Calprotectin is an extremely stable protein, and remains unaltered in stool samples left unprepared for longer than seven days¹³⁷⁻¹³⁹. Studies have focused on the accuracy of faecal calprotectin in the diagnosis and monitoring of IBD. A meta-analysis reported 95% sensitivity and 91% specificity of faecal calprotectin in the identification of IBD. This study also reported that faecal calprotectin was superior to CRP and ESR¹⁴⁰.

The United Kingdom's National Institute for Health and Care Excellence (NICE) recommends faecal calprotectin as a diagnostic tool to help in the differential diagnosis of IBD and IBS¹⁴¹. A normal faecal calprotectin excludes IBD, whereas an elevated faecal calprotectin is an indication for colonoscopy, thereby reducing referrals for unnecessary endoscopic evaluation. A meta-analysis of 13 studies concluded that faecal calprotectin testing would result in a 67% reduction in the number of adults requiring endoscopy, but with a delayed diagnosis in 8% of adults because of false negative results¹⁴².

One area of controversy surrounding faecal calprotectin testing is the determination of an appropriate cut-off value, above which the result is deemed as positive. In most centres, a relatively low level of 50 µg/g is used. A cohort of adult patients undergoing faecal calprotectin testing in primary care was studied¹⁴³. At a cut-off value of 50 µg/g, faecal calprotectin had a negative predictive value (NPV) of 98% and positive predictive value (PPV) of 28%. Increasing the cut-off value to 150 µg/g gave a very comparable NPV of 97%, but a much

higher PPV of 71%. Given these values, it was calculated that increasing the cut-off value to 150 $\mu\text{g/g}$, would reduce colonoscopy and

flexible sigmoidoscopy bookings by 10% at the cost of 4 missed cases of IBD¹⁴³.




Figure 3 – Structure of calprotectin. A three-dimensional model of the crystal structure of Mn^{2+} and Ca^{2+} loaded calprotectin protein, showing two heterodimers: S100A8 and S100A9 as determined by X-ray diffraction. The grey and blue chains represent S100A8 and S100A9, respectively. The purple spheres represent Mn^{2+} and green spheres represent Ca^{2+} . Only one manganese ion can bind per calprotectin heterodimer. The modified figure was taken from the protein data bank (PDB) at: <http://www.rcsb.org/pdb>

12. Blood autophagy genes and nucleotide-binding oligomerization domain-containing protein 2 (NOD2)

Family history of CD is a risk factor for CD¹⁴⁴. Genome-wide association studies have reported that NOD2 and Autophagy genes are associated with CD risk while IL-23/IL-17 is associated with increased risk of both CD and UC¹⁴⁵.

13. Serum anti-neutrophil cytoplasmic antibodies (ANCA) and anti-saccharomyces cerevisiae antibodies (ASCA)

It may be difficult to differentiate CD from UC due to their overlapping pathological, endoscopic and clinical features. Biomarkers like ANCA, which are antibodies against granules of neutrophil cytoplasm and ASCA, which are antibodies against mannan found in the cell walls of *saccharomyces cerevisiae* (*S. cerevisiae*) have been used to help differentiate CD from UC¹⁴⁶.

The sensitivity and specificity of perinuclear ANCA (pANCA) for the diagnosis of ulcerative colitis is 63% and 86% respectively. ASCAs for

diagnosing CD had a sensitivity of 72% and a specificity of 82%¹⁴⁷. The utility of these serological tests in differentiating UC from CD is therefore limited but may be of value in studying disease heterogeneity and disease epidemiology.

Other serologic tests investigated to improve the diagnosis and differential diagnosis of IBD include: antibodies to *Escherichia coli* outer membrane porin C (anti-OmpC), antibodies against laminaribioside (ALCAs), antibodies against chitobioside (ACCAs), *Pseudomonas fluorescens*-associated sequence 12, antibodies to mannoside (AMCAs), pancreatic autoantibodies (PAB), and anti-flagellin CBir1 but all are of limited diagnostic utility.¹⁴⁸⁻¹⁵³

B. Laboratory biomarkers with limited application

The foregoing biomarkers enjoy widespread application in IBD assessment. However, there are some that are not extensively used, or those whose utility for efficacy in IBD assessment can be enhanced only when in combination with one or more biomarkers of similar or different biochemical composition.⁴⁹

These include: adenosine deaminase (ADA), mopterin, nitric oxide, substance P, micro ribonucleic acids (MicroRNAs), lipopolysaccharide-binding protein (LPBP) and cluster of differentiation (CD14), abnormal lectin-based immunoglobulin G (IgG) glycosylation, soluble triggering receptor expressed on myeloid cells-1 (sTREM-1), Soluble suppression of tumorigenicity 2 (ST2), Quantitative fecal immunochemical test (QFIT), activated thrombin activatable fibrinolysis inhibitor (TAFIa), chitinase 3-like-1, mucosal cytokine, mucosal indoleamine 2,3 dioxygenase-1 (IDO1) and angiogenin⁴⁹.

Others are: genes, antibodies toward infliximab, mucosal CHI3L1, urine salicylate level, and thiopurine methyltransferase and 6-thioguanine nucleotide (TPMT)⁴⁹.

14. Inflammatory and serological biomarkers in differential diagnosis of inflammatory bowel disease

The differential diagnosis of IBD includes colon cancer, ischaemic colitis, diverticulitis, bacterial and viral infection. These alternatives or co-existing disorders must be ruled out prior to confirmation of intestinal inflammation as a first step in the differential diagnosis of IBD (*Figure 4*).

The likelihood of CD, UC and Moderate/High Clinical Suspicion are subjected to further radiographic, endoscopic and histological investigations to complete the differential diagnosis. While the above illustration (*figure 4*) provides the necessary information to aid in test selection, interpretation, diagnosis and overall patient management decisions, it is not a substitute for the clinician's underpinning knowledge of IBD in patient assessment based on clinical expertise.¹⁵⁴⁻¹⁵⁷

Accurate diagnosis of CD and UC is therefore dependent on not only laboratory test results but also on the patient's clinical history and examination, histology, imaging results (X-ray, CT and/or MRI scans) and endoscopy^{154,158}.

Inflammation in UC is relatively superficial because it affects the mucosa. It is confined to the colon starting at the rectum, with rare extension to the terminal ileum. There is, however, a 10 to 15% chance of misdiagnosis due to difficulty in distinguishing UC from CD. Accuracy in differential diagnosis is critical because while inflammation in CD extends deeper into the tissues and could affect any portion of the gastrointestinal tract, associated

symptoms like abdominal pain, fever, malnutrition and severe bloody diarrhoea are common to both¹⁵⁹.

15. Inflammatory biomarkers in differential diagnosis of inflammatory bowel disease

The initial step in the identification of IBD is exclusion of other organic disorders. Serum CRP, ESR, faecal calprotectin and faecal lactoferrin form the available inflammatory biomarkers.

Many laboratory biomarkers have been evaluated as the ideal replacement for, or supplement to faecal calprotectin in IBD studies but CRP and ESR remain the most widely used and assessed tests¹. Other biomarkers used include leucocyte counts, platelet counts, albumin levels and orosomucoid concentrations¹⁶⁰, but there is no evidence that they are of greater benefit compared to CRP in the identification of IBD and monitoring its disease activity¹⁶¹.

CRP is routinely available as it is rapid, inexpensive and a simple testing technique. CRP's reliability as a biomarker of choice in routine clinical practice is well established¹⁶². Serum CRP may help in differentiating IBD from other functional bowel disorders¹⁶³. However, normal CRP levels (≤ 5 mg/L) at diagnosis occur in majority of UC patients and in 25% of CD patients.¹⁶⁴ There is, however, a correlation between CRP and endoscopic activity in IBD and this may have a useful role in monitoring responses to therapy.^{165,166}

In summary, increased CRP is more common in CD than in UC, thereby supporting CRP to be a useful biomarker of disease activity in CD than in UC.^{1,164,165} CRP, however, like most serologic biomarkers, is a non-specific biomarker of systemic inflammation.¹

16. Serological biomarkers in differential diagnosis of inflammatory bowel disease

In the presence of inflammation, ANCA and ASCAs may be useful in the identification of IBD.^{154,155} A positive result will necessitate radiology, endoscopy and histological investigation to confirm the diagnosis of IBD and differentiation of CD and UC. If serological markers are negative in the presence of a high clinical suspicion of IBD, then radiologic, endoscopic and histological findings could be used to diagnose or exclude IBD and, if present, to distinguish CD and UC.

ASCAs and atypical pANCA are the two serologic markers most commonly used to distinguish CD and UC.¹⁶⁷ A positive ANCA result is reflexed to determine the relevant pattern(s): cytoplasmic (cANCA), perinuclear (pANCA) or atypical pANCA patterns and their titres. Whereas cANCA and pANCA are found in vasculitis, atypical pANCA is present in IBD. While atypical pANCA is detected in only 5% to 25% of patients with CD, it occurs in about 55% to 80% of those with UC.^{157,168} Conversely, ASCAs is detected in about 60% to 70% of CD patients and only in about 6% to 15% of UC patients.^{168,169}

A combination of pANCA and ASCAs has a sensitivity of 53% and specificity of 93% for both CD and UC; and may be of value in assessing IBD in patients that cannot be distinguished as CD or UC on the basis of indeterminate colitis or established criteria.^{156,157}

Atypical pANCA and ASCAs may help stratify CD. Positive atypical pANCA in CD indicate colonic involvement and an association with a clinical phenotype analogous to UC or UC-like CD, while a positive ASCAs result is linked with the non-UC-like CD.^{169,170} Serologic biomarkers may be of value in children. In children at high risk of IBD, pANCA and

ASCAs may help to identify those children with IBD and to avoid invasive assessment.^{157,171-174}




Figure 4 – The interplay of inflammatory (CRP, ESR, Calprotectin and Lactoferrin) and serological (pANCA and ASCA) biomarkers in the differential diagnosis of inflammatory bowel disease.

Management of disease activity in inflammatory bowel disease

Achieving mucosal healing with subsequent improvement in natural course of the disease state in patients with IBD remains the overall goal of treatment, including the use of immunomodulators and biological agents. Monitoring for efficacy of treatment and relapse

of disease activity, therefore, becomes an important consideration.^{175,176}

No single test procedure or examination has been able to satisfy all the necessary requirements for the clinical management of IBD patients. Assessing disease activity for the foreseeable future involves the use of laboratory tests, radiology, endoscopy, clinical examination and symptoms.¹⁷⁷ This should not, however,

detract from the search for a reliable, non-invasive, highly sensitive and reproducible biomarker of disease activity.^{1,178}

The faecal excretion of ¹¹¹Indium-labelled (¹¹¹In) granulocytes is considered to be the 'gold standard' method for measuring the degree of neutrophilic infiltration into the intestinal mucosa in IBD and hence the disease activity particularly in patients with small bowel CD.^{179,180} The unwieldy nature of radio-labelling techniques in addition to being expensive and involving exposure to radiation restricts their routine clinical use.

Endoscopic assessment of disease activity in IBD, particularly via ileo-colonoscopy is regarded as the 'gold standard' because it has the added advantage of enabling biopsy sampling for histopathological examination apart from offering the opportunity for direct mucosal visualization, endoscopic management of complications, assessing the success of various treatment regimens and predicting the course of disease.¹⁸¹⁻¹⁸⁷ Ileo-colonoscopy, however, is invasive, time-consuming and is limited by expense, risks of complications, patient discomfort and variation in interpretation between endoscopists.^{188,189} This underscores the desirability for an easier method to monitor disease activity.

I. Active versus inactive disease state in inflammatory bowel disease

Increased serum CRP is an indication of active disease in CD patients and it compares favourably with endoscopic disease activity.³⁰ Endoscopic assessment should be considered in patients with UC if elevated serum CRP concentrations fail to normalise with or without underlying symptomology.¹⁹⁰

In the patients with endoscopically confirmed active IBD, faecal calprotectin and lactoferrin

levels correlate with endoscopic disease activity in both CD and UC patients.³⁰ When compared to serum CRP (49%), both faecal calprotectin (88%) and lactoferrin (82%) are more sensitive because of their ability to correlate better with colonic (proctitis: E1 vs. distal colitis: E2 vs. pancolitis: E3) than ileal (ileal and upper: A1 vs. colonic and ileo-colonic: A2) disease activity.^{154,155}

The relatively high sensitivity of both faecal biomarkers (i.e., calprotectin and lactoferrin) in patients with active disease underscores their usefulness in managing patients with IBD¹⁹¹ and thus avoiding endoscopy in patients with a high clinical suspicion of active disease with raised faecal biomarkers. As a corollary, a negative faecal biomarker result may not rule out active disease and therefore endoscopy may be required if clinically indicated.¹⁹¹ Similarly, with CRP, endoscopic assessment should be considered in patients with UC if elevated faecal biomarkers fail to normalise irrespective of underlying symptoms.¹⁹⁰

II. Relapse, remission, and disease course in inflammatory bowel disease

An increase in serum CRP levels predict relapse in disease activity in patients with CD following medically induced remission.^{155,190}

Faecal biomarkers, particularly calprotectin, play a crucial role in predicting relapse in IBD particularly in UC than in CD.¹⁹² Elevated faecal calprotectin level is a common feature in patients with CD who relapse compared to those who remain in remission over a 12-month follow-up.¹⁹³ Low faecal calprotectin levels help identify IBD patients who remain in stable remission during follow-up.¹⁹⁴

Based on a cut-off value of 167 µg/g, faecal calprotectin with a sensitivity of 69% and specificity of 75% appears to be the best

biomarker for predicting IBD relapse following remission.^{154,193} Faecal calprotectin levels may also be particularly valuable in predicting relapse in patients with CD who have undergone surgical resection.^{154,195}

Faecal lactoferrin also appears to have a role in predicting IBD relapse. Elevated faecal lactoferrin levels have a sensitivity of 62% and specificity of 65% in predicting early disease relapse in paediatric patients.^{37,193} In some paediatric patients with CD, pANCA and ASCAs levels also predict complicated disease courses.^{154,155}

III. Response to therapy in inflammatory bowel disease

A normal CRP level in CD patients undergoing treatment is linked to a favourable response to therapy.¹⁹⁰ Whereas normalization of CRP levels may serve as a reliable biomarker to measure the response to therapy, there are currently a paucity of studies investigating the roles of faecal calprotectin, lactoferrin, pANCA and ASCAs in predicting response to therapy.

In summary, more information may be needed on the many aspects (*i.e.*, *analytical and technical factors: sensitivity and specificity, accuracy and precision, reproducibility, calibration, limits of the blank, detection and quantitation; pre-analytical and biological factors: sample handling, patient characteristics, timing, analyte/assay stability; clinical factors and research design variables: purpose, validation, surrogate endpoint, reference ranges*) of laboratory biomarkers to diagnose IBD, differentiate between disease states (CD vs. UC), evaluate disease activity (active vs. quiescent), confirm disease sites or locations as in the small intestine: ileal and upper (A1) vs. colonic and ileo-colonic (A2); large intestine: proctitis (E1) vs. distal colitis (E2) vs. pancolitis (E3), predict the disease

course and relapse, and monitor follow-up and response to treatment.

Summary and future perspectives

It is noteworthy that not all the biomarkers of IBD discovered through clinical research that could be available for routine application in clinical practice. Many of them fail the rigorous protocols of the clinical testing criteria and assay processing through assay development, optimisation and analytical validation, and therefore, are deemed not to be feasible and acceptable for use as laboratory biomarker for assessing IBD. However, for those biomarkers that meet the acceptable target criteria and ‘fitness for purpose’ via the analytical methods, there is also the added laborious intervening timeline of more than 10 years approximation that span from discovery to routine clinical use to consider.^{196,197}

It is a common occurrence for newly discovered biomarkers to fail short of expectation in terms of good analytical performance and diagnostic accuracy during assay validation studies. Such are the demonstrable challenges that are intrinsic in the development and clinical application of biomarkers as articulated in a recent multicentre, open-label randomised controlled trial (RCT) for PRedicting Outcomes For Crohn's disease using a moLecular biomarker (PROFILE).¹⁹⁷

The practical application of laboratory biomarkers in the assessment of inflammatory bowel disease was the subject of a recent study that reviewed the utility of current and potential future biomarkers (*figure 5*).¹⁹⁸ The study reaffirmed the contemporary knowledge of the utility of currently available biomarkers such as CRP and faecal calprotectin to differentiate IBD from functional gastrointestinal pathology and to monitor the disease course. However, it presented such novel biomarkers as Oncostatin M, anti-integrin antibody, faecal MPOs, faecal

microRNAs and glycome profiling, as future biomarkers that could potentially address the risk stratification of individuals not presenting

with the disease (i.e., pre-IBD) including screening those with asymptomatic IBD.¹⁹⁸




Figure 5 – An algorithm for the practical application of laboratory biomarkers in the assessment of inflammatory bowel disease. A narrative of current and potential future biomarkers for diagnosis, prognosis and disease monitoring.¹⁹⁸

The future perspective on the utility of laboratory biomarkers in assessing IBD seems to tilt towards deploying the probable influence of artificial intelligence (AI) to integrate metabolomic, proteomic, genetic and

transcriptomic outputs of biomarker analysis¹⁹⁹⁻²⁰¹. With the continuous growth in complexity of available data, and rapid advances in the availability and affordability of technology, the resort to AI ‘to expand the horizon for biomarker

discovery, enabling the integration of multi-modal data from existing datasets to discover new biomarkers²⁰¹ will be realised.

CONCLUSION

While it is important for laboratory biomarkers of IBD to be non-invasive, cheap, simple, objective, rapid, easy to perform and reproducible, the 'real-world' experience with these biomarkers is increasing together with the increasing number of patients with IBD for which the usefulness of these laboratory biomarkers to assess the disease have become necessary. The 'ideal biomarker' that would integrate all the preceding characteristics for evaluating the analytical performance and diagnostic accuracy of these laboratory biomarkers in IBD is not available. Therefore, the search for the availability of a single laboratory biomarker that would demonstrate the attributes of an 'ideal biomarker' for use in assessing IBD continues.

REFERENCE

1. Vermeire S, van Assche G, Rutgeerts P. Laboratory markers in IBD: useful, magic or unnecessary toys? *Gut*. 2006; 55 (3): 426 – 431.
2. Biomarkers Definitions Working Group. Biomarkers and surrogate endpoints: preferred definitions and conceptual framework. *Clinical Pharmacology and Therapeutics*. 2001; 69 (3): 89 – 95.
3. Lehmann FS, Burri E, Beglinger C. The role and utility of faecal markers in inflammatory bowel disease. *Therapeutic Advances in Gastroenterology*. 2015; 8 (1): 23 – 36.
4. Ballou SP, Kushner I. C-reactive protein and the acute phase response. *Advances in Internal Medicine*. 1992; 37: 313 – 336.
5. Young B, Gleeson M, Cripps AW. C-reactive protein: a critical review. *Pathology*. 1991; 23 (2): 118 – 124.
6. Mold C, Baca R, Du Clos TW. Serum amyloid P component and C-reactive protein opsonise apoptotic cells for phagocytosis through Fc-gamma receptors. *Journal of Autoimmunity*. 2002; 19 (3): 147 – 154.
7. Rudolph WG, Uthoff SM, McAuliffe TL, Goode ET; Petras RE; Galandiuk S. Indeterminate colitis: the real story. *Diseases of the Colon and Rectum*. 2002; 45 (11): 1528 – 1534.
8. Abdelrazeq AS, Wilson TR, Leitch DL, Lund JL, Leveson SH. Ileitis in ulcerative colitis: is it a backwash. *Diseases of the Colon and Rectum*. 2005; 48 (11): 2038 – 2046.
9. Saverymuttu SH, Hodgson HJ, Chadwick VS, Pepsy MB. Differing acute phase responses in Crohn's disease and ulcerative colitis. *Gut*. 1986; 27 (7): 809 – 813.
10. Sidoroff M, Karikoski R, Raivio T, Savilahti E; Kolho KL. High-sensitivity C-reactive protein in paediatric inflammatory bowel disease. *World Journal of Gastroenterology*. 2010; 16 (23): 2901 – 2906.
11. Yamamoto-Furusho JK, Camacho-Escobedo J, Tellez-Avilla F, Barreto R. β 2-microglobulin levels and high-sensitive C-reactive protein as markers of histological activity in patients with chronic idiopathic ulcerative colitis. *Gaceta Medica de Mexico*. 2010; 146 (1): 31 – 35.
12. Gross V, Andus T, Caesar I, Roth M, Schölmerich J. Evidence for continuous stimulation of interleukin-6 production in

Crohn's disease. *Gastroenterology*. 1992; 102 (2): 514 – 519.

13. Szalai AJ, McCrory MA, Cooper GS, Wu J, Kimberly RP. Association between baseline levels of C-reactive proteins (CRP) and a dinucleotide repeat polymorphism in intron of the CRP gene. *Genes Immunology*. 2002; 3 (1): 14 – 19.

14. Russell AI, Cunninghame-Graham DS, Shepherd C, Robertson CA, Whittaker J, Meeks J *et al*. Polymorphism at the C-reactive protein locus influences gene expression and predisposes to systemic lupus erythematosus. *Human Molecular Genetics*, 2004; 13 (1): 137 – 147.

15. Carlson CS, Aldred SF, Lee PK, Tracy RP, Schwartz SM, Rieder M *et al*. Polymorphism with the C-reactive protein (CRP) promoter region are associated with plasma CRP levels. *American Journal of Human Genetics*. 2005; 77 (1): 64 – 77.

16. Willot S, Vermeire S, Ohresser M, Rutgeerts P, Paintaud G, Belaiche J *et al*. C-reactive protein gene polymorphisms are not associated with biological or clinical response to infliximab in Crohn's disease. *Gastroenterology*. 2005; 128 (Suppl.): A311

17. Danese J, Wheeler JG, Hirschfield GM, Eda S, Eiriksdottir G, Rumley A *et al*. C-reactive protein and other circulating markers of inflammation in the prediction of coronary heart disease. *New England Journal of Medicine*. 2004; 350 (14): 1387 – 1397.

18. Collins CE, Rampton DS. Platelet dysfunction: a new dimension in inflammatory bowel disease. *Gut*. 1995; 36 (1): 5 – 8.

19. Dong WG, Liu SP, Zhu HH, Luo HS, Yu JP. Abnormal function of platelets and role of

angelica sinensis in patients with ulcerative colitis. *World Journal of Gastroenterology*. 2004; 10 (4): 606 – 609.

20. Danese S, de la Motte C, Fiocchi C. Platelets in inflammatory bowel disease: clinical, pathogenic and therapeutic implications. *The American Journal of Gastroenterology*. 2004; 99 (5): 938 – 945.

21. Kapsoritakis AN, Koukourakis MI, Sfiridaki A, Potamianos SP, Kosmadaki MG, Koutroubakis IE *et al*. Mean platelet volume: a useful marker of inflammatory bowel disease activity. *The American Journal of Gastroenterology*. 2001; 96 (3): 776 – 781.

22. Kayahan H, Akarsu M, Ozcan MA, Demir S Ates H, Unsal B *et al*. Reticulated platelet levels in patients with ulcerative colitis. *International Journal of Colorectal Disease*. 2007; 22 (12): 1429 – 1435.

23. Zubcevic N, Mesihovic R, Zubcevic S. Usefulness of laboratory data in estimation of Crohn's disease activity. *Medicinski Arhiv.*, 2010; 64 (1): 33 – 36.

24. Song CS, Park DI, Yoon MY, Seok HS, Park JH, Kim HJ *et al*. Association between red cell distribution with and disease activity in patients with inflammatory bowel disease. *Digestive Diseases and Sciences*. 2012; 57 (4): 1033 – 1038.

25. Baveye S, Ellass E, Mazurier J, Spik G, Legrand D. Lactoferrin: a multifunctional glycoprotein involved in the modulation of the inflammatory process. *Clinical Chemistry and Laboratory Medicine*. 1999; 37 (3): 281 – 286.

26. Kane SV, Sandborn WJ, Rufo PA, Zholudev A, Boone J, Lyerly D *et al*. Faecal lactoferrin is a sensitive and specific marker in identifying

intestinal inflammation. *The American Journal of Gastroenterology*. 2003; 98 (6): 1309 – 1314.

27. Desai D, Faubion WA, Sandborn WJ. Biological activity markers in inflammatory bowel disease. *Alimentary Pharmacology and Therapeutics*. 2007; 25 (3): 247 – 255.

28. Joishy M, Davies L, Ahmed M, Wassel J, Davies K, Sayers A *et al*. Faecal calprotectin and lactoferrin as noninvasive markers of pediatric inflammatory bowel disease. *Journal of Pediatric Gastroenterology and Nutrition*. 2009; 48 (1): 48 – 54.

29. Sugi K, Saitoh O, Hirata I, Katsu K. Faecal lactoferrin as a marker of disease activity in inflammatory bowel disease: comparison with other neutrophil-derived proteins. *The American Journal of Gastroenterology*. 1996; 91 (5): 927 – 934.

30. Jones J, Loftus EV Jr., Panaccione R, Chen LS, Peterson S, McConnell J *et al*. Relationships between disease activity and serum and faecal biomarkers in patients with Crohn's disease. *Clinical Gastroenterology & Hepatology*. 2008; 6 (11): 1218 – 1224.

31. Walker TR, Land ML, Kartashov A, Saslowsky TM, Lyerly DM, Boone JH *et al*. Faecal lactoferrin is a sensitive and specific marker of disease activity in children and young adults with inflammatory bowel disease. *Journal of Paediatric Gastroenterology Nutrition*. 2007; 44 (4): 414 – 422.

32. Dai J, Liu WZ, Zhao YP, Hu YB; Ge ZZ. Relationship between faecal lactoferrin and inflammatory bowel disease. *Scandinavian Journal of Gastroenterology*. 2007; 42 (12): 1440 – 1444.

33. Sipponen T, Karkkainen P, Savilahti E, Kolho K, Nuutinen H, Turunen U *et al*.

Correlation of faecal calprotectin and lactoferrin with an endoscopic score for Crohn's disease and histological findings. *Alimentary Pharmacology Therapeutics*. 2008; 28 (10): 1221 – 1229.

34. Schoepfer A, Trummler M, Seeholzer P, Seibold-Schmid B; Seibold F. Discriminating IBD from IBS: comparison of the test performance of faecal markers, blood leucocytes, CRP and IBD antibodies. *Inflammatory Bowel Disease*. 2008; 14 (1): 32 – 39.

35. Langhorst J, Elsenbruch S, Koelzer J, Rueffer A, Michalsen A, Dobos GJ. Non-invasive markers in the assessment of intestinal inflammation in inflammatory bowel disease: performance of faecal lactoferrin, calprotectin and PMN-elastase, CRP and clinical indices. *The American Journal of Gastroenterology*. 2008; 103 (1): 162 – 169.

36. Sidhu R, Wilson P, Wright A, Yau CW, D'Cruz FA, Foye L *et al*. Faecal lactoferrin – a novel test to differentiate between the irritable and inflamed bowel? *Alimentary Pharmacology Therapeutics*. 2010; 31 (12): 1365 – 1370.

37. Gisbert J, McNicholl A, Gomollon F. Questions and answers on the role of faecal lactoferrin as a biological marker in inflammatory bowel disease. *Inflammatory Bowel Disease*. 2009; 15 (11): 1746 – 1754.

38. D'Inca R, Dal Pont E, Di Leo V, Ferronato A, Fries W, Vettorato MG *et al*. Calprotectin and lactoferrin in the assessment of intestinal inflammation and organic disease. *International Journal of Colorectal Diseases*. 2007; 22 (4): 429 – 437.

39. Moishy M, Davies I, Ahmed M, Wassel J, Davies K, Sayers A *et al*. Faecal calprotectin and lactoferrin as non-invasive markers of paediatric inflammatory bowel disease. *Journal of*

Paediatric Gastroenterology and Nutrition. 2009; 48 (1): 48 – 54.

40. Nancey S, Boschetti G, Moussata D, Cotte E, Peyras J, Cuerq C *et al.* Neopterin is a novel reliable faecal marker as accurate as calprotectin for predicting endoscopic disease activity in patients with inflammatory bowel diseases. *Inflammatory Bowel Diseases*. 2013; 19 (5): 1043 – 1052.

41. Husain N, Tokoro K, Popov JM, Naides SJ, Kwasny MJ, Buchman AL *et al.* Neopterin concentration as an index of disease activity in Crohn's disease and ulcerative colitis. *Journal of Clinical Gastroenterology*. 2013; 47 (3): 246 – 251.

42. Annaházi A, Molnár T, Farkas K, Rosztóczy A, Izbéki F, Gecse K *et al.* Faecal MMP-9: a new noninvasive differential diagnostic and activity marker in ulcerative colitis. *Inflammatory Bowel Disease*. 2013; 19 (2): 316 – 320.

43. Kolho K, Sipponen T, Valtonen E, Savilahti E. Faecal calprotectin, MMP-9 and human beta-defensin-2 levels in paediatric inflammatory bowel disease. *International Journal of Colorectal Diseases*. 2014; 29 (1): 43 – 50.

44. Silberer H, Küppers B, Mickisch O, Baniewicz W, Drescher M, Traber L *et al.* Faecal leucocyte proteins in inflammatory bowel disease and irritable bowel syndrome. *Clinical Laboratory*. 2005; 51 (3-4): 117 – 126.

45. Schroeder O, Naumann M, Shastri Y, Povse N; Stein J. Prospective evaluation of faecal neutrophil-derived proteins in identifying intestinal inflammation: combination of parameters does not improve diagnostic accuracy of calprotectin. *Alimentary Pharmacology Therapeutics*. 2007; 26 (7): 1035 – 1042.

46. Czub E, Herzig KH, Szaflarska-Popawska, Kiehne K, Socha P, Woś H *et al.* Faecal pyruvate kinase: a potential new marker for intestinal inflammation in children with inflammatory bowel disease. *Scandinavian Journal of Gastroenterology*. 2007; 42 (10): 1147 – 1150.

47. Turner D, Leach ST, Mack D, Uusoue K, McLernon R, Hyams J *et al.* Faecal calprotectin, lactoferrin, M2-pyruvate kinase and S100A12 in severe ulcerative colitis: a prospective multicentre comparison of predicting outcomes and monitoring response. *Gut*. 2010; 59 (9): 1207 – 1212.

48. Chung-Faye G, Hayee B, Maestranzi S, Donaldson N; Forgacs I; Sherwood R. Faecal M2-pyruvate kinase (M2-PK): a novel marker of intestinal inflammation. *Inflammatory Bowel Disease*. 2007; 13 (11): 1374 – 1378.

49. Fengming Y, Jianbing W. Biomarkers of inflammatory bowel disease. *Disease Markers*. 2014;710915. Doi:10.1155/2014/710915.

50. Pietzsch J, Hoppmann S. Human S100A12: a novel key player in inflammation? *Amino Acids*. 2009; 36 (3): 381 – 389.

51. Schafer BW, Heizmann CW. The S100 family of EF-hand calcium-binding proteins: functions and pathology. *Trends in Biochemical Science*. 1996; 21 (4): 134 – 140.

52. Heizmann CW; Cox JA. New perspectives on S100 proteins: a multifunctional Ca²⁺-, Zn²⁺- and Cu²⁺-binding protein family. *BioMetals*. 1998; 11 (4): 383 – 397.

53. Donato R. Intracellular and extracellular roles of S100 proteins. *Microscopy Research & Technique*. 2003; 60 (6): 540 – 551.

54. Vogl T, Propper C, Harmann M, Strey A, Strupat K, van den Bos C *et al.* S100A12 is

expressed exclusively by granulocytes and acts independently from MRP8 and MRP14. *Journal of Biological Chemistry*. 1999; 274 (36): 25291 – 25296.

55. Karl J, Wild N, Tacke M, Andres H, Garczarek U, Rollinger W *et al*. Improved diagnosis of colorectal cancer using a combination of faecal occult blood and novel faecal protein markers. *Clinical Gastroenterology & Hepatology*. 2008; 6 (10): 1122 – 1228.

56. Foell D, Wittkowski H, Ren Z, Turton J, Pang G, Daebritz J *et al*. Phagocyte-specific S100 proteins are released from affected mucosa and promote immune responses during inflammatory bowel disease. *Journal of Pathology*. 2008; 216 (2): 183 – 192.

57. Ravasi T, Hsu K, Goyette J, Schroder K, Yang Z, Rahimi F *et al*. Probing the S100 protein family through genomic and functional analysis. *Genomics*. 2004; 84 (1): 10 – 22.

58. Dell'Angelica EC, Schleicher CH, Santome JA. Primary structure and binding properties of calgranulin C, a novel S100-like calcium-binding protein from pig granulocytes. *Journal of Biological Chemistry*. 1994; 269 (46): 28929 – 28936.

59. Guignard F, Mauel J, Markert M. Identification and characterization of a novel human neutrophil protein related to the S100 family. *Biochemical Journal*. 1995; 309 (Pt 2): 395 – 401.

60. Fritz G, Botelho HM, Morozova-Roche LA, Gomes CM. Natural and amyloid self-assembly of S100 proteins: structural basis of functional diversity. *FEBS Journal*. 2010; 277:4578 – 4590.

61. Santamaria-Kisiel L, Rintala-Dempsey AC, Shaw GS. Calcium-dependent and -independent interactions of the S100 protein family. *Biochemical Journal*. 2006; 396 (2): 201 – 214.

62. Donato R. S100: a multigenic family of calcium-modulated proteins of the EF-hand type with intracellular and extracellular functional roles. *International Journal of Biochemistry and Cell Biology*, 2001; 33 (7): 637 – 668.

63. Heizmann CW, Fritz G, Schafer B. S100 proteins: structure, functions and pathology. *Frontiers in Bioscience: A Journal and Virtual Library*. 2002; 7: d1356 – 1368.

64. Ye F, Foell D, Hirono KI, Vogl T, Rui C, Yu X *et al*. Neutrophil-derived S100A12 is profoundly upregulated in the early stage of acute Kawasaki disease. *American Journal of Cardiology*. 2004; 94 (6): 840 – 844.

65. Donato R, Cannon BR, Sorci G *et al*. Functions of S100 Proteins. *Current Molecular Medicine*. 2013; 13 (1): 24 – 57.

66. Manolakis AC, Kapsoritakis AN, Tiaka EK, Potamianos SP. Calprotectin, calgranulin C and other members of the S100 protein family in inflammatory bowel disease. *Digestive Disease Science*. 2011; 56 (6): 1601 – 1611.

67. Hoffmann MA, Drury S, Fu C, Qu W, Taguchi A, Lu Y *et al*. RAGE mediates a novel proinflammatory axis: a central cell surface receptor for S100/calgranulin polypeptides. *Cell*, 1999; 97 (7): 889 – 901.

68. Hitomi J, Yamaguchi K, Kikuchi Y, Kimura T, Maruyama K, Nagasaki K. A novel calcium-binding protein in amniotic fluid, CAAF-1: its molecular cloning and tissue distribution. *Journal of Cell Science*. 1996; 109 (Pt 4): 805 – 815.

69. Schmidt AM, Yan SD, Yan SF, Stern DM. The biology of the receptor for advanced glycation end products and its ligands. *Biochimica et Biophysica Acta*. 2000; 1498 (2–3): 99 – 111.
70. Foell D, Frosch M, Sorg C, Roth J. Phagocyte-specific calcium-binding S100 proteins as clinical laboratory markers of inflammation. *Clinica Chimica Acta*. 2004a. 344 (1–2): 37 – 51.
71. Foell D, Wittkowski H, Hammerschmidt I, Wulffraat N, Schmeling H, Frosch M *et al*. Monitoring neutrophil activation in juvenile rheumatoid arthritis by S100A12 serum concentrations. *Arthritis & Rheumatology*. 2004b. 50 (4): 1286 – 1295.
72. Yang Z, Tao T, Raftery MJ, Youssef P, Di Girolamo N, Geczy CL. Proinflammatory properties of the human S100 protein S100A12. *Journal of Leukocyte Biology*. 2001; 69 (6): 986 – 994.
73. Rouleau P, Vandal K, Ryckman C, Poubelle PE, Boivin A, Talbot M. The calcium-binding protein S100A12 induces neutrophil adhesion, migration, and release from bone marrow in mouse at concentrations similar to those found in human inflammatory arthritis. *Clinical Immunology*. 2003; 107 (1): 46 – 54.
74. Foell D, Kane D, Bresnihan B, Vogl T, Nacken W, Sorg C *et al*. Expression of the pro-inflammatory protein S100A12 (EN-RAGE) in rheumatoid and psoriatic arthritis. *Rheumatology (Oxford)*. 2003a. 42 (11): 1383 – 1389.
75. Liao H, Wu J, Kuhn E, Chin W, Chang B, Jones MD *et al*. Use of mass spectrometry to identify protein biomarkers of disease severity in the synovial fluid and serum of patients with rheumatoid arthritis. *Arthritis & Rheumatology*. 2004; 50 (12): 3792 – 3803.
76. Kosaki A, Hasegawa T, Kimura T, Lida K, Hitomi J, Matsubara H *et al*. Increased plasma S100A12 (EN-RAGE) levels in patients with type 2 diabetes. *Journal of Clinical Endocrinology & Metabolism*. 2004; 89 (11): 5423 – 5428.
77. Nakashima M, Sakai T, Hiraiwa H, Hamada T, Omachi T, Ono Y *et al*. Role of S100A12 in the pathogenesis of osteoarthritis. *Biochemical and Biophysical Research Communications*. 2012; 422 (3): 508 – 514.
78. Thierolf M, Hagmann M-L, Pfeffer M, Berntenis N, Wild N, Roeßler M *et al*. Towards a comprehensive proteome of normal and malignant human colon tissue by 2-D-LC-ESI-MS and 2-DE proteomics and identification of S100A12 as a potential cancer biomarker. *Proteomics Clinical Application*. 2008; 2 (1): 11 – 22.
79. Kallinich T, Wittkowski H, Keitzer R, Roth J, Foell D. Neutrophil-derived S100A12 as a novel biomarker of inflammation in familial mediterranean fever. *Annals of Rheumatic Diseases*. 2010; 69 (4): 677 – 682.
80. Miniati M, Monti S, Basta G, Cocci F, Fornai E, Bottai M. Soluble receptor for advanced glycation end products in COPD: relationship with emphysema and chronic cor pulmonale: a case-control study. *Respiratory Research*. 2011; 12: 37
81. Hofmann-Bowman M, Wilk J, Heydemann A, Kim G, Rehman J, Lodato JA *et al*. S100A12 mediates aortic wall remodelling and aortic aneurysm. *Circulation Research*. 2008; 106 (1): 145 – 154.
82. Mori Y, Kosaki A, Kishimoto N, Kimura T, Lida K, Fukui M *et al*. Increased plasma S100A12 (EN-RAGE) levels in haemodialysis

patients with atherosclerosis. *American Journal of Nephrology*. 2009; 29 (1): 18 – 24.

83. Madland TM, Larsen A, Brun JG. S100 proteins calprotectin and S100A12 are related to radiographic changes rather than disease activity in psoriatic arthritis with low disease activity. *Journal of Rheumatology*. 2007; 34 (10): 2089 – 2092.

84. Kalousova M, Kubena AA, Benakova H, Dusilova-Sulkova S, Tesar V, Zima T. ENRAGE (extracellular newly identified receptor for advanced glycation end-products binding protein) and mortality of long-term haemodialysis patients: a prospective observational cohort study. *Clinical Biochemistry*. 2012; 45 (7–8): 556 – 560.

85. Shiotsu Y, Mori Y, Nishimura M, Sakoda C, Tokoro T, Hatta T *et al*. Plasma S100A12 level is associated with cardiovascular disease in haemodialysis patients. *Clinical Journal of American Society of Nephrology*. 2011; 6 (4): 718 – 723.

86. Kikkawa T, Sato N, Kojika M, Takahashi G, Aoki K, Hoshikawa K *et al*. Significance of measuring S100A12 and sRAGE in the serum of sepsis patients with postoperative acute lung injury. *Digestive Surgery*. 2010; 27 (4): 307 – 312.

87. Wittkowski H, Sturrock A, van Zoelen MAD, Viemann D, van der Poll T, Hoidal JR *et al*. Neutrophil-derived S100A12 in acute lung injury and respiratory distress syndrome. *Critical Care Medicine*. 2007; 35 (5): 1369 – 1375.

88. Hermani A, Hess J, De Servi B, Medunjanin S, Grobholz R, Trojan L *et al*. Calcium-binding proteins S100A8 and S100A9 as novel diagnostic markers in human prostate cancer. *Clinical Research: An Official Journal of the*

American Association for Cancer Research. 2005; 11 (14): 5146 – 5152.

89. Sidler MA, Leach ST, Day AS. Faecal S100A12 and faecal calprotectin as innovative markers for inflammatory bowel disease in children. *Inflammatory Bowel Disease*. 2008; 14 (3): 359 – 366.

90. de Jong Naomi SH, Leach ST, Day AS. Faecal S100A12: A novel non-invasive marker in children with Crohn's disease. *Inflammatory Bowel Disease*. 2006; 12 (12): 566 – 572.

91. Moroz OV, Antson AA, Murshudov GN, Maitland NJ, Dodson GG, Wilson KS *et al*. The three-dimensional structure of human S100A12. *Acta Crystallographica. Section D, Biological Crystallography*, 2001; 57 (Pt. 1): 20 – 29.

92. Moroz OV, Antson AA, Dodson EJ, Burrell HJ, Grist SJ, Lloyd RM *et al*. The structure of S100A12 in a hexameric form and its proposed role in receptor signalling. *Acta Crystallographica. Section D, Biological Crystallography*, 2002; 58 (Pt. 3): 407– 413.

93. Moroz OV, Dodson GG, Wilson KS, Lukanidin E, Bronstein IB. Multiple structural states of S100A12: a key to its functional diversity. *Microscopic Research and Technique*, 2003; 60 (6): 581 – 592.

94. Moroz OV, Antson AA, Grist SJ, Maitland NJ, Dodson GG, Wilson KS *et al*. Structure of the human S100A12-copper complex: implications for host-parasite defence. *Acta Crystallographica. Section D, Biological Crystallography*, 2003; 59 (Pt. 5): 859 – 867.

95. Fritz G, Botelho HM, Morozova-Roche LA, Gomes CM. Natural and amyloid self-assembly of S100 proteins: structural basis of functional diversity. *FEBS Journal*, 2010; 277: 4578 – 4590.

96. Fagerhol MK; Dale I; Andersson T. A radioimmunoassay for a granulocyte protein as a marker in studies on the turnover of such cells. *Bulletin of European Physiopathology and Respiration*, 1980; 16 Suppl: 273 – 282.
97. Dale I, Brandtzaeg P, Fagerhol MK, Scott H. Distribution of a new myelomonocytic antigen (L1) in human peripheral blood leukocytes. Immunofluorescence and immunoperoxidase staining features in comparison with lysozyme and lactoferrin. *American Journal of Clinical Pathology*, 1985; 84 (1): 24 – 34.
98. Brandtzaeg P, Dale I, Fagerhol MK. Distribution of a formalin-resistant myelomonocytic antigen (L1) in human tissues. I. Comparison with other leukocyte markers by paired immunofluorescence and immunoenzyme staining. *American Journal of Clinical Pathology*, 1987; 87 (6): 681 – 699.
99. Brandtzaeg P, Dale I, Fagerhol MK. Distribution of a formalin-resistant myelomonocytic antigen (L1) in human tissues. II. Normal and aberrant occurrence in various epithelia. *American Journal of Clinical Pathology*, 1987; 87 (6): 700 – 707.
100. Dale I, Brandtzaeg P. Expression of the epithelial L1 antigen as an immunohistochemical marker of squamous cell carcinoma of the lung. *Histopathology*, 1989; 14 (5): 493 – 502.
101. Berntzen HB, Fagerhol MK. L1, a major granulocyte protein; isolation of high quantities of its subunits. *Scandinavian Journal of Clinical Laboratory Investigation*, 1990; 50 (7): 769 – 774.
102. Berntzen HB, Olmez U, Fagerhol MK, Munthe E. The leukocyte protein L1 in plasma and synovial fluid from patients with rheumatoid arthritis and osteoarthritis. *Scandinavian Journal of Rheumatology*, 1991; 20 (2): 74 – 82.
103. Berntzen HB, Fagerhol MK, Ostensen M, Mowinckel P; Hoyeraal HM. The L1 protein as a new indicator of inflammatory activity in patients with juvenile rheumatoid arthritis. *Journal of Rheumatology*, 1991; 18 (1): 133 – 138.
104. Dale I. Plasma levels of the calcium-binding L1 leukocyte protein: standardization of blood collection and evaluation of reference intervals in healthy controls. *Scandinavian Journal of Clinical & Laboratory Investigation*, 1990; 50 (8): 837 – 841.
105. Santhanagopalan V, Hahn BL, Dunn BE, Weissner JH; Sohnle PG. Antimicrobial activity of calprotectin isolated from human empyema fluid supernatants. *Clinical Immunology and Immunopathology*, 1995; 76 (3 Pt 1): 285 – 290.
106. Steinbakk M, Naess-Andresen CF, Lingaas E, Dale I, Brandtzaeg P, Fagerhol MK. Antimicrobial actions of calcium binding leucocyte L1 protein, calprotectin. *Lancet*, 1990; 336 (8718): 763 – 765.
107. Johne B, Fagerhol MK, Lyberg T, Prydz H, Brandtzaeg P, Naess-Andresen CF. Functional and clinical aspects of the myelomonocyte protein calprotectin. *Molecular Pathology (MP)*, 1997; 50 (3): 113 – 123.
108. Sampson B, Fagerhol MK, Sunderkötter C, Golden BE, Richmond P, Klein N et al. Hyperzincaemia and hypercalprotectinaemia: a new disorder of zinc metabolism. *Lancet*, 2002; 360 (9347): 1742 – 1745.
109. Kerkhoff C, Vogl T, Nacken W, Sopalla C, Sorg C. Zinc-binding reverses the calcium-induced arachidonic acid-binding capacity of

- the S100A8/A9 protein complex. *FEBS Letters*, 1999; 460 (1): 134 – 138.
110. Isaksen B, Fagerhol MK. Calprotectin inhibits matrix metalloproteinases by sequestration of zinc. *Molecular Pathology*, 2001; 54 (5): 289 – 292.
111. Sohnle PG, Hunter MJ, Hahn B, Chazin WJ. Zinc-reversible antimicrobial activity of recombinant calprotectin (migration inhibitory factor-related proteins 8 and 14). *Journal of Infectious Diseases*, 2000; 182 (4): 1272 – 1275.
112. Tibble JA, Bjarnason I. Fecal calprotectin as an index of intestinal inflammation. *Drugs Today (Barc)*, 2001; 37 (2): 85 – 96.
113. Larsen A, Hovdenak N, Karlsdottir A, Wentzel-Larsen T, Dahl O, Fagerhol MK. Faecal calprotectin and lactoferrin as markers of acute radiation proctitis: a pilot study of eight stool markers. *Scandinavian Journal of Gastroenterology*, 2004; 39 (11): 1113 – 1118.
114. Healy AM, Pickard MD, Pradhan AD, Wang Y, Chen Z, Croce K *et al.* Platelet expression profiling and clinical validation of myeloid-related protein 14 as a novel determinant of cardiovascular events. *Circulation*, 2006; 113 (19): 2278 – 2284.
115. Altwegg LA, Neidhart M, Hersberger M, Müller S, Eberli FR, Corti R *et al.* Myeloid-related protein 8/14 complex is increased by monocytes and granulocytes at the site of coronary occlusion: a novel, early and sensitive marker of acute coronary syndromes. *European Heart Journal*, 2007; 28 (8): 941 – 948.
116. Wilkinson MM, Busuttill A, Hayward C, Brock DJ, Dorin JR, Van Heyningen V. Expression pattern of two related cystic fibrosis-associated calcium-binding proteins in normal and abnormal tissues. *Journal of Cell Science*, 1988; 91 (Pt 2): 221 – 230.
117. Yui S, Nakatani Y, Mikami M. Calprotectin (S100A8/S100A9), an inflammatory protein complex from neutrophils with a broad apoptosis-inducing activity. *Biological & Pharmaceutical Bulletin*, 2003; 26 (6): 753 – 760.
118. Striz I; Trebichavsky I. Calprotectin – a pleiotropic molecule in acute and chronic inflammation. *Physiological Research / Academia Scientiarum Bohemoslovaca*, 2004; 53 (3): 245 – 253.
119. Müller F, Froland SS, Aukrust P, Fagerhol MK. Elevated serum calprotectin levels in HIV-infected patients: the calprotectin response during ZDV treatment is associated with clinical events. *Journal of Acquired Immune Deficiency Syndromes*, 1994; 7 (9): 931 – 939.
120. Madland TM, Hordvik M, Haga HJ, Jonsson R, Brun JG. Leukocyte protein calprotectin and outcome in rheumatoid arthritis. *Scandinavian Journal of Rheumatology*, 2002; 31 (6): 351 – 354.
121. Berntzen HB, Munthe E, Fagerhol MK. A longitudinal study of the leukocyte protein L1 as an indicator of disease activity in patients with rheumatoid arthritis. *Journal of Rheumatology*, 1989; 16 (11): 1416 – 1420.
122. Brun JG, Haga HJ, Bøe E, Kallay I, Lekven C, Berntzen HB *et al.* Calprotectin in patients with rheumatoid arthritis: relation to clinical and laboratory variables of disease activity. *The Journal of Rheumatology*, 1992; 19 (6): 859 – 862.
123. Hammer HB, Kvien TK, Glennas A, Melby K. A longitudinal study of calprotectin as an inflammatory marker in patients with reactive

arthritis. *Clinical & Experimental Rheumatology*, 1995; 13 (1): 59 – 64.

124. Wittkowski H, Frosch M, Wulffraat N, Goldbach–Mansky R, Kallinich T, Kuemmerle–Deschner J *et al.* S100A12 is a novel molecular marker differentiating systemic onset juvenile idiopathic arthritis from other causes of fever of unknown origin. *Arthritis & Rheumatology*, 2008; 58 (12): 3924 – 3931.

125. Brun JG, Jonsson R, Haga HJ. Measurement of plasma calprotectin as an indicator of arthritis and disease activity in patients with inflammatory rheumatic diseases. *The Journal of Rheumatology*, 1994; 21 (4): 733 – 738.

126. Kane D, Roth J, Frosch M, Vogl T, Bresnihan B, FitzGerald O. Increased perivascular synovial membrane expression of myeloid–related proteins in psoriatic arthritis. *Arthritis Rheumatology*, 2003; 48 (6): 1676 – 1685.

127. Brun JG, Madland TM, Gran JT, Myklebust G. A longitudinal study of calprotectin in patients with polymyalgia rheumatica or temporal arteritis: relation to disease activity. *Scandinavian Journal of Rheumatology*, 2005; 34 (2): 125 – 128.

128. Haga HJ, Brun JG, Berntzen HB, Cervera R, Khamashta M, Hughes GR. Calprotectin in patients with systemic lupus erythematosus: relation to clinical and laboratory parameters of disease activity. *Lupus*, 1993; 2 (1): 47 – 50.

129. Burkhardt K, Radespiel–Tröger M, Rupperecht HD, Goppelt–Struebe M, Riess R, Renders L *et al.* An increase in myeloid–related protein serum levels precedes acute renal allograft rejection. *Journal of American Society of Nephrology*, 2001; 12 (9): 1947 – 1957.

130. Stríz I, Jaresová M, Lácha J, Sedláček J, Vítko S. MRP8/14 and pro– calcitonin serum levels in organ transplantations. *Annals of Transplantation*, 2001; 6 (2): 6 – 9.

131. Heller F, Frischmann S, Grumbaum M, Zidek W, Westhoff TH. Urinary calprotectin and the distinction between prerenal and intrinsic acute kidney injury. *Clinical Journal of the American Society of Nephrology: CJASN*, 2011; 6 (10): 2347 – 2355.

132. Lönnkvist MH, Theodorsson E, Holst M, Ljung T, Hellstrom PM. Blood chemistry markers for evaluation of inflammatory activity in Crohn’s disease during infliximab therapy. *Scandinavian Journal of Gastroenterology*, 2011; 46 (4): 420 – 427.

133. Börekçi B, Aksoy H, Ozturk N, Kadanali S. Correlation between calprotectin and oxidised LDL in pre–eclampsia. *Turkish Journal of Medical Sciences*, 2009; 39 (2): 191 – 195.

134. Røseth AG, Aadland E, Jahnsen J, Raknerud N. Assessment of disease activity in ulcerative colitis by faecal calprotectin, a novel granulocyte marker protein. *Digestion*, 1997; 58 (2): 176 – 180.

135. Røseth AG, Schmidt PN, Fagerhol MK. Correlation between faecal excretion of indium–111–labelled granulocytes and calprotectin, a granulocyte marker protein, in patients with inflammatory bowel disease. *Scandinavian Journal of Gastroenterology*, 1999; 34 (1): 50 – 54.

136. Røseth AG. Determination of faecal calprotectin, a novel marker of organic gastrointestinal disorders. *Digestive Liver Disease*, 2003; 35 (9): 607 – 609.

137. Guardiola J, Lobatón T, Cerrillo E, Ferreiro–Iglesias R, Gisbert JP, Domènech E *et al.* Recommendations of the Spanish Working Group on Crohn's Disease and Ulcerative Colitis (GETECCU) on the utility of the determination of faecal calprotectin in inflammatory bowel

disease. *Gastroenterology y Hepatologia*, 2018; 41 (8): 514 – 529.

138. Røseth A, Fagerhol M, Aadland E, Schjønby H. Assessment of the neutrophil dominating protein calprotectin in faeces. A methodological study. *Scandinavian Journal of Gastroenterology*, 1992; 27 (9): 793 – 798.

139. Tøn H, Brandsnes Ø, Dale S, Holtlund J, Skuibina E, Schjønby H *et al.* Improved assay for faecal calprotectin. *Clin Chim Acta*, 2000; 292 (1 – 2): 41 – 54.

140. von Roon AC, Karamountzos L, Purkayastha S, Reese GE, Darzi AW, Teare JP *et al.* Diagnostic precision of fecal calprotectin for inflammatory bowel disease and colorectal malignancy. *American Journal of Gastroenterology*, 2007; 102 (4): 803 – 813.

141. National Institute for Health and Care Excellence. Faecal calprotectin diagnostic tests for inflammatory diseases of the bowel. *NICE Diagnostics Guidance*, 2013; 11: 1 – 58.

142. van Rheene P, Van de Vijver E, Fidler V. Faecal calprotectin for screening of patients with suspected inflammatory bowel disease: diagnostic meta-analysis. *British Medical Journal*, 2010; 341: c3369.

143. Pavlidis P, Chedgy FJ, Tibble JA. Diagnostic accuracy and clinical application of faecal calprotectin in adult patients presenting with gastrointestinal symptoms in primary care. *Scandinavian Journal of Gastroenterology*, 2013; 48 (9): 1048 – 1054.

144. Khor B, Gardet A, Xavier RJ. Genetics and pathogenesis of inflammatory bowel disease. *Nature*, 2011; 474 (7351): 307 – 317.

145. Cho H, Brant SR. Recent insights into the genetics of inflammatory bowel disease. *Gastroenterology*, 2011; 140 (6): 1704 – 1712.

146. Ferrante M, Henckaerts L, Joossens M, Pierik M, Joossens S, Dotan N *et al.* New serological markers in inflammatory bowel

disease are associated with complicated disease behaviour. *Gut*, 2007; 56 (10): 1394 – 1403.

147. Linskens RK, Mallant-Hent RC, Groothuisink ZMA, Bakker-Jonges LE, van de Merwe JP, Hooijkaas, H *et al.* Evaluation of serological markers to differentiate between ulcerative colitis and Crohn's disease: pANCA, ASCA and agglutinating antibodies to anaerobic coccoid rods. *European Journal of Gastroenterology & Hepatology*, 2002; 14 (9): 1013 – 1018.

148. Prideaux L, de Cruz P, Ng SC, Kamm MA. Serological antibodies in inflammatory bowel disease: a systematic review. *Inflammatory Bowel Disease*, 2012; 18 (7): 1340 – 1355.

149. Lakatos PL, Papp M, Rieder F. Serologic antiglycan antibodies in inflammatory bowel disease. *The American Journal of Gastroenterology*, 2011; 106 (3): 406 – 412.

150. Bogdanos DP, Rigopoulou EI, Smyk DS, Roggenbuck D, Reinhold D, Forbes A *et al.* Diagnostic value, clinical utility and pathogenic significance of reactivity to the molecular targets of Crohn's disease specific– pancreatic autoantibodies. *Autoimmunity Reviews*, 2011; 11 (2): 143 – 148.

151. Zholudev A, Zurakowski D, Young W, Leichtner A, Bousvaros A. Serologic testing with ANCA, ASCA and anti-OmpC in children and young adults with Crohn's disease and ulcerative colitis: diagnostic value and correlation with disease phenotype. *American Journal of Gastroenterology*, 2004; 99 (11): 2235 – 2241.

152. Dotan I, Fishman S, Dgani Y, Schwartz M, Karban A, Lerner A *et al.* Antibodies against laminaribioside and chitobioside are novel serologic markers in Crohn's disease. *Gastroenterology*, 2006; 131 (2): 366 – 378.

153. Mendoza JL, Abreu MT. Biological markers in inflammatory bowel disease: practical consideration for clinicians.

Gastroenterologie Clinique et Biologie, 2009; 33 (Suppl. 3): S158 – S173.

154. Iskandar HN, Ciorba MA. Biomarkers in inflammatory bowel disease: current practices and recent advances. *Translational Research*, 2012; 159 (4): 313 – 325.

155. Lewis JD. The utility of biomarkers in the diagnosis and therapy of inflammatory bowel disease. *Gastroenterology*, 2011; 140 (6): 1817 – 1826.

156. Kornbluth A, Sachar DB. Practice Parameters Committee of the American College of Gastroenterology. Ulcerative colitis practice guidelines in adult: American College of Gastroenterology, Practice Parameters Committee. *American Journal of Gastroenterology*, 2010; 105 (3): 501 – 523; quiz 524.

157. Reese GE, Constantinides VA, Simillis C, Darzi AW, Orchard TR, Fazio VW *et al.* Diagnostic precision of anti-saccharomyces cerevisiae antibodies and perinuclear antineutrophil cytoplasmic antibodies in inflammatory bowel disease. *American Journal of Gastroenterology*, 2006; 101 (10): 2410 – 2422.

158. Soubières AA, Poullis A. Emerging role of novel biomarkers in the diagnosis of inflammatory bowel disease. *World Journal of Gastrointestinal Pharmacology and Therapeutics*, 2016; 7 (1): 41 – 50.

159. Savige J, Dimech W, Fritzler M, Goeken J, Hagen EC, Jennette JC *et al.* Addendum to the International Consensus Statement on testing and reporting of antineutrophil cytoplasmic antibodies. Quality control guidelines, comments, and recommendations for testing in other autoimmune diseases. International Group for Consensus Statement on Testing and Reporting of Antineutrophil Cytoplasmic Antibodies (ANCA). *American Journal of Clinical Pathology*, 2003; 120 (3): 312 – 318.

160. Cioffi M, De Rosa A, Serao R, Picone I, Vietri MT. Laboratory markers in ulcerative colitis: current insights and future advances. *World Journal of Gastrointestinal Pathophysiology*, 2015; 6 (1): 13 – 22.

161. Vilela EG, Torres HO, Martins FP, Ferrari Mde L, Andrade MM, Cunha AS. Evaluation of inflammatory activity in Crohn's disease and ulcerative colitis. *World Journal of Gastroenterology*, 2012; 18 (9): 872 – 881.

162. Vermeire S, Van Assche G, Rutgeerts P. C-reactive protein as a marker for inflammatory bowel disease. *Inflammatory Bowel Disease*, 2004; 10 (5): 661 – 665.

163. Shine B, Berghouse L, Jones JE, Landon J. C-reactive protein as an aid in the differentiation of functional and inflammatory bowel disorders. *Clinica Chimica Acta.*, 1985; 148 (2): 105 – 109.

164. Henriksen M, Jahnsen J, Lygren I, Stray N, Sauar J, Vatn MH *et al.* C-reactive protein: A predictive factor and marker of inflammation in inflammatory bowel disease. Results from a prospective population-based study. *Gut*, 2008; 57 (11): 1518 – 1523.

165. Solem CA, Loftus EV Jr., Tremaine WJ, Harmsen WS, Zinsmeister AR, Sandborn WJ *et al.* Correlation of C-reactive protein with clinical, radiological and endoscopic activity in inflammatory bowel disease (IBD). *Inflammatory Bowel Disease*, 2005; 11 (8): 707 – 712.

166. Reinisch W, Wang Y, Oddens BJ, Link R. C-reactive protein, an indicator for maintained response or remission to infliximab in patients with Crohn's disease: a post-hoc analysis from ACCENT I. *Alimentary Pharmacology & Therapeutics*, 2012; 35 (5): 568 – 576.

167. Zhou F, Xia B, Wang F, Shrestha UK, Chem M, Wang H *et al.* The prevalence and diagnostic value of perinuclear antineutrophil cytoplasmic antibodies and anti-saccharomyces cerevisiae antibodies in patients with

inflammatory bowel disease in mainland China. *Clinica Chimica Acta.*, 2010; 411 (19 – 20): 1461 – 1465.

168. Bossuyt X. Serologic markers in inflammatory bowel disease. *Clinical Chemistry*, 2006; 52 (2): 171 – 181.

169. Abreu MT, Vasiliauskas EA, Kam LY, Dubinsky MC. Use of serologic tests in Crohn's disease. *Clinical Perspectives in Gastroenterology*, 2001; 4 (3): 155 – 164.

170. Klebl FH, Bataille F, Berthea CR, Herfarth H, Hofstädter F, Schölmerich J *et al.* Association of perinuclear antineutrophil cytoplasmic antibodies and anti-saccharomyces cerevisiae antibodies with Vienna classification subtypes of Crohn's disease. *Inflammatory Bowel Disease*, 2003; 9 (5): 302 – 307.

171. Viennois E, Baker MT, Xiao B, Wang L, Laroui H, Merlin D. Longitudinal study of circulating protein biomarkers in inflammatory bowel disease. *Journal of Proteomics*, 2015; 112: 166 – 179.

172. Van der Vijvers E, Schreuder A, Cnossen W, Muller Kobold A, van Rheenen P. Safely ruling out inflammatory bowel disease in children and teenagers without referral for endoscopy. The North Netherlands Paediatric IBD Consortium. *Archives of Disease of Childhood*, 2012; 97 (12): 1014 – 1018.

173. Bartůnková J, Kolárová I, Sedivá A, Hölzelová E. Antineutrophil cytoplasmic antibodies, anti-saccharomyces cerevisiae antibodies, and specific IgE to food allergens in children with inflammatory bowel diseases. *Clinical Immunology*, 2002; 102 (2): 162 – 168.

174. Ruemmele FM, Targan SR, Levy G, Dubinsky M, Braun J, Seidman EG. Diagnostic accuracy of serological assays in paediatric inflammatory bowel disease. *Gastroenterology*, 1998; 115 (4): 822 – 829.

175. De vos M, Dewit O, D'Haens G, Baert F, Fontaine F, Vermeire S *et al.* Fast and sharp decrease in calprotectin predicts remission by infliximab in anti-TNF naïve patients with ulcerative colitis. *Journal of Crohn's and Colitis*, 2012; 6 (5): 557 – 562.

176. Dignass A, Van Assche G, Lindsay JO, Lémann M, Söderholm J, Colombel JF *et al.* European Crohn's and Colitis Organisation (ECCO) The second European evidence-based consensus on the diagnosis and management of Crohn's disease: current management. *Journal of Crohn's and Colitis*, 2010; 4 (1): 28 – 62.

177. Soubières AA, Poullis A. Emerging role of novel biomarkers in the diagnosis of inflammatory bowel disease. *World Journal of Gastrointestinal Pharmacology and Therapeutics*, 2016; 7 (1): 41 – 50.

178. Vrabie R, Kane S. Non-invasive markers of disease activity in inflammatory bowel disease. *Gastroenterology & Hepatology*, 2014; 10 (9): 576 – 584.

179. Bjarnason I, MacPherson A, Hollander D. Intestinal permeability: an overview. *Gastroenterology*, 1995; 108 (5): 1566 – 1581.

180. Saverymuttu SH, Peters AM, Lavender JP, Hodgson HJ, Chadwick VS. ¹¹¹Indium autologous leucocytes in inflammatory bowel disease. *Gut*, 1983; 24 (4): 293 – 299.

181. Matsuura M, Saito D, Hisamatsu T. The role of endoscopy in the diagnosis and treatment of IBD. *The Japanese Journal of Gastroenterology*, 2021; 118 (7): 607 – 617.

182. Spiceland CM, Lodhia N. Endoscopy in inflammatory bowel disease: role in diagnosis, management and treatment. *World Journal of Gastroenterology*, 2018; 24 (35): 4014 – 4020.

183. Chang S, Malter L, Hudesman D. Disease monitoring in inflammatory bowel disease. *World Journal of Gastroenterology*, 2015; 21 (40): 11246 – 11259.

184. Carter D, Eliakim R. Current role of endoscopy in inflammatory bowel disease diagnosis and management. *Current Opinion in Gastroenterology*, 2014; 30 (4): 370 – 377.
185. Carter D, Lang A, Eliakim R. Endoscopy in inflammatory bowel disease. *Minerva Gastroenterologica e Diabetologica*, 2013; 59 (3): 273 – 284.
186. Dignass A, Eliakim R, Magro F, Maaser C, Chowers Y, Geboes K *et al.* Second European evidence-based consensus on the diagnosis and management of ulcerative colitis Part 1: Definitions and diagnosis. *Journal of Crohn's and Colitis*, 2012; 6 (10): 965 – 990.
187. Van Assche G, Dignass A, Panes J, Beaugerie L, Karagiannis J, Allez M *et al.* The second European evidence-based consensus on the diagnosis and management of Crohn's disease: definitions and diagnosis. *Journal of Crohn's and Colitis*, 2010; 4 (1): 7 – 27.
188. Mitselos IV, Fousekis FS, Lamouri C, Katsanos KH, Christodoulou DK. Current noninvasive modalities in Crohn's disease monitoring. *Annals of Gastroenterology*, 2021; 34 (6): 770 – 780.
189. Zhou FS, Gao N, Sun X, Jiang XY, Chen JJ, Mao QQ *et al.* C-reactive protein/albumin ratio is a useful biomarker for predicting the mucosal healing in the Crohn's disease: A retrospective study. *Medicine (Baltimore)*, 2021; 100 (10): e24925. Doi:10.1097/MD.000000000024925.
190. Peyrin-Biroulet L, Sandborn W, Sands BE, Reinisch W, Bemelman W, Bryant RV *et al.* Selecting therapeutic targets in inflammatory bowel disease (STRIDE): Determining therapeutic goals for treat-to-target. *American Journal of Gastroenterology*, 2015; 110 (9): 1324 – 1338.
191. Mosli MH, Zou G, Garg SK, Feagan SG, MacDonald JK, Chande N *et al.* C-reactive protein, fecal calprotectin and stool lactoferrin for detection of endoscopic activity in symptomatic inflammatory bowel disease patients: a systematic review and meta-analysis. *American Journal of Gastroenterology*, 2015; 110 (6): 802 – 819.
192. Costa F, Mumolo MG, Ceccarelli L, Bellini M, Romano MR, Sterpi C *et al.* Calprotectin is a stronger predictive marker of relapse in ulcerative colitis than in Crohn's disease. *Gut*, 2005; 54 (3): 364 – 368.
193. Gisbert JP, Bermejo F, Pérez-Calle JL, Taxonera C, Vera I, McNicholl AG *et al.* Fecal calprotectin and lactoferrin for the prediction of inflammatory bowel disease relapse. *Inflammatory Bowel Disease*, 2009; 15 (8): 1190 – 1198.
194. Mooiweer E, Severs M, Schipper ME, Fidler HH, Siersema PD, Laheji RJ *et al.* Low fecal calprotectin predicts sustained clinical remission in inflammatory bowel disease patients: a plea for deep remission. *Journal of Crohn's and Colitis*, 2015; 9 (1): 50 – 55.
195. D'Inca R, Dal Pont E, Di Leo V, Benazzato L, Martinato M, Lamboglia F *et al.* Can calprotectin predict relapse risk in inflammatory bowel disease? *American Journal of Gastroenterology*, 2008; 103 (8): 2007 – 2014.
196. Armbruster DA, Pry T. Limit of blank, limit of detection and limit of quantitation. *Clinical Biochemistry Review*, 2008; 29 (Suppl. 1): S49 – S52.
197. Noor NM, Lee JC, Bond S, Dowling F, Brezina B, Pavel KV *et al.* A biomarker-stratified comparison of top-down versus accelerated step-up treatment strategies for patients with newly diagnosed Crohn's disease (PROFILE): a multicentre, open-label randomised controlled trial. *The Lancet. Gastroenterology & Hepatology*, 2024; 9 (5): 415 – 427.
198. Clough J; Colwill M; Poullis A, Pollock R, Patel K, Honap S. Biomarkers in inflammatory

bowel disease: a practical guide. *Therapeutic Advances in Gastroenterology*, 2024; 17: 1 – 19. DOI: 10.1177/17562848241251600

199. Mao R, Chen M. Precision medicine in IBD: genes, drugs, bugs and omics. *Nature Reviews. Gastroenterology & Hepatology*, 2022; 19 (2): 81 – 82.

200. Atreya R, Neurath MF. Biomarkers for personalizing IBD therapy: the quest continues. *Clinical Gastroenterology and Hepatology: The*

Official Clinical Practice Journal of the American Gastroenterological Association, 2024; 22 (7): 1353 – 1364. DOI: 10.1016/j.cgh.2024.01.026

201. Prelaj A, Miskovic V, Zanitti M, Trovo F, Genova C, Viscardi G et al. Artificial intelligence for predictive biomarker discovery in immuno-oncology: a systematic review. *Annals of Oncology: Official Journal of the European Society for Medical Oncology*, 2024; 35 (1): 29 –65.