

Toxicological studies of ethanol leaf extract of *Lavandulastoechas* on kidney of Wistar rat.Joseph Oyepata Simeon*¹, Joseph Opeyemi Tosin²¹Department of Pharmacology and Toxicology, Faculty of Pharmaceutical Sciences, Federal University, Oye-Ekiti, Ekiti State, Nigeria.²Department of Pharmacology, Faculty of Pharmacy, Lead City University, Ibadan, Oyo State, NigeriaSubmitted: 9th Jan., 2023; Accepted: 9th March., 2023; Published: 30th April., 2023DOI: <https://doi.org/10.54117/jcbr.v3i2.11>*Corresponding author: Joseph Oyepata Simeon; oyepata.joseph@fuoye.edu.ng; simeon4unme@yahoo.com**Abstract**

The medicinal ingredients in herbal remedies are derived from plants. Despite the fact that they could offer a wide range of health advantages, they are thought to have a lower likelihood for adverse effects. The medicinal herb lavender (*Lavandulastoechas* L.) is widely utilized for a variety of medical conditions. The aim of this investigation is to evaluate the toxicological effect of *Lavandulastoechas* on Wistar rat kidney. Twenty four (24) Animals of either sex were used. Group 1 was given distilled water (10 ml/kg), while groups 2, 3, and 4 were given 50, 100, and 200 mg/kg of *Lavandulastoechas*, respectively for 28 days. Animals were housed in typical cages and given oral access to the extract for 28 days before being weighed and put to death. A heart puncture was used to get blood, which was then promptly taken for testing. Red blood cell, Hemoglobin, and Mean corpuscular volume levels decreased significantly ($P < 0.05$), but neutrophils, basophiles, eosinophils, and platelets remained constant. At 100 and 200 mg/kg, *Lavandulastoechas* significantly ($P < 0.05$) lowered K^+ levels. Na^+

increased significantly ($P < 0.05$) at a dosage of 200 mg/kg. Across doses given, the levels of creatinine, chloride, and urea were not substantially ($P < 0.05$) impacted. A histological analysis shows minimal tubular deformation. The study's findings indicated that the plant might have a negligible impact on the kidney, indicating that it should be used with caution if consumed over an extended period of time.

Key: *Lavandulastoechas*, blood, rats, kidney**Introduction**

Modern medicine has its roots in the ancient therapeutic practices that have been practiced long before recorded human history. Numerous common medications, including morphine (opium poppies), digoxin (foxglove), quinine (cinchona bark), and aspirin (willow bark), were developed using plant materials (Solomon et al., 2015). The usage of natural medicines has dramatically expanded across the globe. Since prehistoric times, medicinal plants have been discovered and used in traditional medical procedures. Many chemical compounds are produced by plants for a variety of purposes, including

defense against herbivorous mammals, fungus, insects, and illnesses (Ojochegebe et al., 2019; Joseph and Joseph 2018). Different phytochemicals with potential or established biological activity have been identified (Joseph et al., 2018; Timothy et al., 2018). The results of taking a complete plant as medication remain unclear, though, because a single plant includes a vast variety of phytochemicals. Furthermore, extensive scientific research to determine the phytochemical composition and pharmacological effects of many plants with therapeutic promise is still lacking (Aprioku et al., 2016; Okokon et al., 2017). In non-industrialized communities, medicinal plants are frequently employed, mostly because they are more accessible and affordable than modern treatment (Gaitonde, 2017). Testing for sub-acute toxicity is used to evaluate toxicity in the target species. It provides insight into the relative toxicity of compounds, allowing them to be categorized as mild, moderate, very hazardous, and almost non-toxic. Additionally, it offers details on the process by which drugs exert their hazardous effects and offers a foundation for selecting doses for efficacy studies as well as long-term toxicity research (Joseph et al., 2020; Simeon et al., 2020).

Subacute toxicity test is defined by the globally harmonized system (GHS) as specific target organ/system toxicity resulting from 28 days of recurrent chemical exposure in rodents (Builder et al., 2019; Joseph et al., 2019). Testing for sub-acute toxicity is performed to assess impacts primarily on different organ systems and to determine a no observed effect level (Eknoyan et al., 1997;

Haruna et al., 2020). The evaluation of clinical observation, blood analysis, a full body gross necropsy, and a microscopic examination of all organs and tissues make up the end points (Wazis et al., 2020; Builder et al., 2020). Herbal plants have been used for many years for therapeutic and health purposes. Local people frequently ingest medicinal plants without following a prescribed dosage schedule or time frame (Zubairu et al., 2021; Joseph et al., 2021). This may have unintended consequences for the tissue, organ, or biological system (Nahon et al., 2011).

Although there is little to no scientific evidence about the efficacy and safety of medicinal plants, long-term clinical experience has generally served as the basis for their use (Romer and Parsons 1977). The use of medicinal herbs as a remedy is only based on a long-standing, widespread traditional folk practice (Romer and Parsons 1977; Oluwakanyesola et al., 2018). A thorough scientific examination of these plants is essential given the rise in the use of herbal medicines and the necessity to support their traditional uses (Tosin et al., 2019; Simeon et al., 2019). Although herbs are supposed to be safe, there have been many reports of unsafe and fatal side effects (Emamian et al., 1993; Molina and DiMaio 2015). These could include overt toxicity, allergic responses, contaminant effects, and/or medicine and herb interactions. Because they are "natural," phytotherapeutic treatments are frequently wrongly thought to be less hazardous. However, those items have bioactive components that could have negative impacts (Boron 2004).

Various aesthetic and therapeutic uses for lavender (*Lavandula stoechas*), a plant belonging to the Labiatae family, are found in herbal medicine (Simeon et al., 2019; Sabastine et al., 2019; Joseph et al., 2019). *Lavandulastoechas* essential oil inhalation decreased cholesterol plaques in rabbits with atherosclerotic disease but had no impact on serum cholesterol levels (Samson et al., 2019). In rats, lavender had a hypolipidemic impact (Simeon et al., 2019). Lavender aromatherapy has additionally shown vasodilatory benefits and increased coronary blood flow in people (Joseph et al., 2019). Isolated rat hearts were guarded against ischemia reperfusion (IR) damage by a lavender flower extract (Oyebadejo et al., 2019). In an experimental model of stroke, lavender oil had neuroprotective efficacy and antioxidant properties (Joseph et al., 2019b). In a recently completed study, therapy with lavender essential oil after myocardial infarction decreased ischemia injury in rats (Modupe et al., 2019). Several studies have been carried out on this plant but no report on safety or sub chronic toxicity study has been carried out. The aim of this investigation is to evaluate the toxicological effect of *Lavandulastoechas* on Wistar rat kidney.

Material and Method

Wistar rats, both sexes, were acquired from Bingham University's Animal House. They were fed typical animal pellets and given unlimited amounts of water. The College of Health Sciences Animal Ethics Committee at Bingham University granted permission and consent for the use of animals in research as

BU/DR&D/REC/019/003. It plant was also given the voucher number BUFP 3001.

Plant collection

In the adjacent Karu village in Nasarawa State, Nigeria, leaves from the *Lavandulastoechas* plant were harvested from its natural environment in the morning during raining season. The Department of Botany at Bingham University in Nasarawa State, Nigeria, verified the plant's authenticity. It was given the voucher number BUFP 3001.

Plant extraction

For two weeks, the leaves were dried in the shadows. The ground-up dried plant material was further broken up and pulverized. The substance was pulverized and then macerated in 70% ethanol. Utilizing a rotary evaporator in a vacuum, the liquid filtrates were concentrated and evaporated to dryness at 40°C. Until it was used, the ethanol extract was kept at -4°C.

Animal study

Twenty four rats of either sex (weighing 174–257g) were chosen at random and divided into four groups of six rats each. Rats in groups 2, 3, and 4 got 50, 100, and 200 mg/kg of *Lavandulastoechas* extract, respectively, while group 1 acted as the control group and received normal saline (10ml/kg). At the start of the experiment and once a week, the weights of the rats were noted. The day of

sacrifice was given the designation D₂₉, whereas the first day of dosing was given the designation D₀.

Haematological analysis

On the day 29, of the trial, the rats were sacrificed and blood was collected through Cardiac puncture. EDTA-containing sample bottles were used to collect a portion of the blood for hematological analysis, which included measuring hemoglobin concentration, white blood cell counts (WBC), differentials (neutrophils, eosinophils, basophils, lymphocytes, and monocytes), red blood cell counts (RBC), platelets, and hemoglobin (Hb) concentration (Cell-Dyn, Abbott, USA).

Chempathology analysis

Second portion of the blood was collected into plain bottle, allowed to clot and centrifuged at 300rpm for 10 minutes. The serum collected was used to estimate biochemical parameters.

Histological study

The kidney of the animals were surgically removed and weighed and a part of each was

fixed in 10% formaldehyde for histological processes.

Statistical analysis

Data were expressed as the Mean ±Standard Error of the Mean (SEM). Data were analyzed statistically using one-way Analysis of Variance (ANOVA) followed by Dunnett’s post hoc test for multiple comparisons between the control and treated groups. Values of P ≤ 0.05 were considered significant.

Results

Effect of 28 days oral administration of *Lavandulastoechason* hematological parameters in rats

At a dose of 100 mg/kg, *Lavandulastoechas* considerably (p<0.05) decreased the levels of red blood cells, hemoglobin, and platelets; however, at a dose of 50 mg/kg, it significantly (p<0.05) increased the mean corpuscular hemoglobin concentration in rats compared to the control. However, the mean corpuscular hemoglobin concentration did not substantially (p<0.05) impact the level of basophiles, neutrophils, eosinophils, or lymphocytes. (Table 1).

Table 1: Effect of 28 days oral administration of ethanol leaf extract of *LavandulaLavandulastoechason* hematological parameters in wistar rats.

Hematological parameters	DW(10ml/kg)	50	Treatment (mg/kg)	
			100	200
WBC (×10 ⁹ /L)	8.27±0.77	6.54±1.42	3.72±0.67*	7.20±1.83
RBC (×10 ¹² /L)	8.30±0.36	8.65±0.64	6.17±0.25*	7.74±0.25
HGB (g/dL)	15.90±0.56	15.24±0.63	11.36±0.77*	14.58±0.36

HCT (g/dL)	55.18±2.02	56.61±3.56	34.67±3.29*	53.40±1.80
MCV (fL)	66.67±0.94	65.44±1.45	57.17±0.10*	69.60±1.72
MCH (pg)	19.16±0.16	17.80±1.19	18.83±0.38	18.80±0.22
MCHC (g/dL)	29.15±0.16	27.43±1.23	32.51±0.61*	27.10±0.67
PLT (×10 ⁹ /L)	620.83±52.81	567.00±96.28	252.00±50.14*	670.45±55.78*
LYM (%)	86.81±4.61	85.00±4.03	82.83±5.82	86.41±3.14
NEUT (×10 ⁹ /L)	10.81±3.64	10.82±3.61	15.40±5.41	11.00±3.23
EOSI (×10 ⁹ /L)	1.50±0.33	2.40±0.48	1.80±0.44	1.25±0.22
BASO (×10 ⁹ /L)	1.00±0.29	2.00±0.55	2.50±1.51	3.30±2.21

Data presents Mean ± SEM: n = 6, One way ANOVA, followed by Dunnett's post hoc for multiple comparison

*significantly different from the distilled water (DW) control at p<0.05. DW = distilled water (WBC = white blood cells, RBC = red blood cells, HGB = hemoglobin, HCT = hematocrit, MCV = mean corpuscular volume, MCH = mean corpuscular hemoglobin, MCHC = mean corpuscular hemoglobin concentration, PLT = platelet, LYM = lymphocyte, NEUT = neutrophils, EOSI = eosinophils, BASO = basophils).

Effect of 28 days oral administration of *Lavandulastoechason* renal indices and electrolytes in Wistar rats.

At 100 and 200 mg/kg, *Lavandulastoechas* significantly (p<0.05) lowered K⁺ levels. Na⁺ increased significantly (p<0.05) at a dosage of 200 mg/kg. Across doses given, the levels of creatinine, chloride, and urea were not significantly affected (p<0.05).

Table 4.4: Effect of 28 days oral administration of ethanol leaf extract of *Lavandulastoechason* renal indices in Wister rats.







Renal indices and electrolytes (mmol/L)	Treatment (mg/kg)			
	DW(10ml/kg)	LS (50)	LS (100)	LS (200)
Potassium	7.00±0.55	7.60±0.70	5.30±0.48*	5.8±0.20*

Sodium	158.00±2.90	168.20±2.91	169.10±1.90*	164.20±1.79
Chloride	101.30±5.81	99.72±6.61	111.23±2.46	100.55±3.21
Urea	9.43±0.29	9.54±0.55	8.90±0.25	8.75±0.53
Creatinine	63.44±9.62	80.22±12.41*	55.11±16.932	73.71±6.11

Data presents Mean ± SEM: n = 6, *significantly different from the distilled water (DW) control at p <0.05. DW = distilled water.

Histopathological Investigations of the effect of 28 days oral administration of *Lavandulastoechason* renal indices and electrolytes in Wistar rats.

At 100 mg/kg and 200 mg/kg, the kidney exhibited mild tubular deformation and glomerular necrosis. Additionally, there was mild tubular necrosis along with lymphocyte hyperplasia at 50 mg/kg dose. Histology characteristics of Kidney of rats in the control group were normal.






Plate XV: Histological sections of Kidneys of rats treated with Normal saline 10 ml/kg bw (1), LS 50 mg/kg bw (2), LS 100 mg/kg bw (3) and LS 200 mg/kg bw (4) at magnification A (x100) and B(x400) Stained with H&E technique.

Keys: Renal corpuscle (RC) Convoluted tubules (CT), Collecting ducts (CD), Epithelial lining degeneration (ELD), Pyknotic nucleus (Pn) Glomerulus (GL), Nucleus (N) Tubular necrosis (TN) and vascular degeneration (VD)




Fig 1: graph showing effect of ethanol leaf extract of *L. On* rats' serum sodium, chloride and potassium levels. X axis represent doses administered while Y axis represent serum level of sodium, chloride and potassium.

Discussion

This study assessed how the plant affected the kidney after administering on rats for a period of time. When compared to the control group of rats, the ethanol extract of *Lavandula stoechas* caused a substantial decrease in the red blood cell, hemoglobin, and platelet counts. This suggested that the plant might affect the process and production of red blood cells, by shorten their life span, or affect how the body consumes iron. Anemia, or low red blood cell count, can make a person feel weak and worn out. The body has to work harder to supply enough oxygen to the cells when the red blood count is lower than normal. A low red blood cell (RBC) count can result in a wide range of symptoms and medical issues (Sabastine et al., 2021; Tosin et al., 2021). The blood's hemoglobin helps carry oxygen from your lungs to your tissues. In muscle cells, myoglobin takes up, stores, transports, and releases oxygen (Oyepata 2021). Additionally, the extract had no effect on the amount of basophiles, neutrophils, eosinophils, or lymphocytes. This suggests that the plant might not have an impact on the immune system

The study also demonstrated that while other parameters remained mostly unchanged, *Lavandula stoechas* slightly increased serum potassium ions and decreased serum sodium ions. This could suggest that the ethanol leaf extract of the plant does not affect electrolyte level when consumed for a sustained period. The extract had no effect on creatinine levels,

indicating that it may have no toxicological or therapeutic significant for causing or treating kidney-related disorders. According to several studies (Weinstein 1994; Joseph et al.; Kalantar-Zadeh et al., 2021; Kalantar-Zadeh et al., 2021), Some kidney diseases can cause an increase in creatinine level due to loss of the creatinine's normal excretory function, damage to the muscles, or after taking an unsuitable medication that interferes with the kidney's normal function (Simeon et al., 2022; Novick et al., 2006). Creatinine is mostly produced by tissue creatinine breakdown from endogenous sources (Thomas 2005). Also, in this study the level of Serum urea was unchanged when compared to the control group. Serum urea concentration is frequently regarded as a more accurate indicator of renal function than serum creatinine. Serum/plasma urea concentration reflects the balance between urea production in the liver and urea elimination by the kidneys. An increased in plasma/serum urea can be caused by increased urea production, decreased urea elimination which affects the glomerular filtration rate, or a combination of the two (Oyepata and Simeon 2022; Maton et al., 1993). Result of the study suggests that the extract does not increase protein breakdown and has little significant on glomerular filtration rate. Hence, ethanol leaf extract of *Lavandula stoechas* does not affect the kidney functions.

The kidneys of the rats treated with *Lavandula stoechas* displayed slight

pathological symptoms of injury, including degraded microvesicles in the tubular lining cells among others, according to the histological results. Kidney sections of the control group rats examined showed normal glomeruli that were free of pathological symptoms. This agrees with other results previous work (Weinstein 1994; Joseph et al.; Kalantar-Zadeh et al., 2021; Kalantar-Zadeh et al., 2021) and in that the extract may not possess severe toxicological significant when consumed for a particular period of time

Conclusion

The histological examination and biochemical parameters results revealed that *Lavandulastoechas* has chemical constituents that may somewhat modify the anatomical and physiological characteristics of the kidney, indicating that care should be taken while consuming it over an extended period of time.

Acknowledgement

The author wishes to thank everyone who has contributed to the success of this research work.

Conflict of interest

Author declares that there is no conflict of interest.

References

Adhvaryu M, Reddy N, Parabia MH. (2007). Effects of four Indian medicinal herbs on Isoniazid -, Rifampicin - and Pyrazinamide -

induced hepatic injury and immunosuppression in guinea pigs. *World J Gastroenterol.*; 13(23): 3199 – 3205.

Ahn K. (2017). The worldwide trend of using botanical drugs and strategies for developing global drugs. *BMB Reports.* **50** (3): 111–116.

Aprioku JS, Joseph OS, Obianime AW (2016). Quantification of Antinociceptive and Anti-Inflammatory Potentials of Different *Ocimum gratissimum* Linn. Leaf Extracts in Whistar Albino Rats. *European Journal of Medicinal Plants.* Volume 17(3). Page 1-8.

Bhuiyan MD, Mohashine JG, Nishimura M, Matsumura ME, Seishi T, Shimono T. (1997). Antibacterial effects of the crude *Azaadirachta indica*. Neem bark extract of streptococcus sobrinus. *Pediatric Dental Journal.*, 7(1);61-64.

Bhuiyan MD, Mohashine JG, Nishimura M, Matsumura ME, Seishi T, Shimono T. (1997). Antibacterial effects of the crude *Azaadirachta indica*. Neem bark extract of streptococcus sobrinus. *Pediatric Dental Journal.* 7(1);61-64.

Builder MI, Joseph SO, Olugbemi TO, Akande, T (2020). Toxicity. Studies of extract of African Mistletoe: *Agelanthus Dodoneifolius* Polh and Wiens in Rats. *Nigeria biomedical Journal.* Page 113-130
Builder MI, Anzaku SA, Joseph SO (2019). Effectiveness of intermittent preventive treatment in pregnancy with sulphadoxine-pyrimethamine against malaria in northern Nigeria. *International Journal of Recent*

Scientific Research Vol. 10 (05), pp. 32295-32299.

Builders MI, Joseph SO, Bassi PU. (2020). A Survey of Wound Care Practices by Nurses in a Clinical Setting. International Journal of Healthcare and Medical Sciences. Vol. 6, Issue. 5, Page 74-81.

Builders MI, Simeon JO, Ogundeko TO, Builders P. (2020). Antimalarial Drugs and COVID -19. Sumerianz Journal of Medical and Healthcare. Vol. 3, No. 12, pp. 111-116.

Edeoga HO, Okwe DE, Mbabie BO Phytochemical constituents of some Nigerian Medicinal plant. Agric_JBiotechnol. 2005, 4:685-688

Endo TY, Haraguch, K, Saka WD. Renal toxicity in rats after oral administration of mercury contaminated boiled whole livers marked for human consumption .Archives of Environmental Contamination and Toxicology .Journal of Pharmaceutical Research, 2003, 44(3):p.412-416.

Haruna WC, Builders M, Simeon JO, Tosin JO (2020). Toxicological Study of the Effect of Ethanol Leaf Extract of *Pterocarpuss antalinus* Extract on Liver of Wister Rats. Nigeria biomedical Science Journal. Page 17-29.

Haruna WC, Simeon JO, Builders M, Tosin JO (2020). Effect of ethanol leaf extract of *cassia angustifolia* extract on heartand lipid profile of wister rats. African Journal of Pharmaceutical Research & Development. Vol. 12 No.1. Page 1-8.

Izzo A. (2004). Drug interactions with St. John's Wort (*Hypericum perforatum*): A review of the clinical evidence. International Journal of Clinical Pharmacology and Therapeutics; 42:139-148.

Joseph OS, Musa TL, Joseph O T, Ibhafidon I. (2020). The Dynamics of Differential Impacts of COVID-19 on African Countries Compared to Other Parts of the World. International journal of multidisciplinary research and analysis. Volume 03 Issue 11. Page 185-198.

Joseph OS, Sabastine AZ, Joseph OT, Adegbuyi TA. (2022). An Analysis of Daily distributive effect of COVID-19 Pandemic across the Globe Using the USA as a Comparism Factor: An update report of 17th of February, 2022.

Joseph OS, Joseph OT. (2018). Hepatoprotective activity of ethanol stem extract of *Homalium letestui* against thioacetamide-induced liver injury. The Nigerian Journal of Pharmacy. Vol. 52 (1). Page 67-74.

Joseph OS, Builders M, Joseph OT. (2020). Effect of Caffeine on Diazepam - Induced Sedation and Hypnosis in Wister Rat. Global Scientific Journal. Vol. 8, Issue 9. Page 451-466

Joseph OS, Builders M, Joseph OT, Sabastine AZ. (2020). Assessing differential impacts of COVID-19 on African countries: A comparative study. International Journal of Research and Innovation in Applied Science. Vol. 5, Issue 5. Page 197-203

- Joseph OS, Joseph OT, Musa TL, Oyepata PJ. (2019). Histological evaluation of the nephroprotective activity of the ethanol stem extracts of *Homalium letestui* in Gentamicin – induced albino rats injury, using various staining techniques. Global Scientific Journal. Volume 7, Issue 8. Page 1065-1087.
- Joseph OS, Jude EO, Joseph OT. (2018). Hepatoprotective activity of extract of *Homalium letestui* stem against carbon tetrachloride-induced liver injury. Advance Herbal Medicine. Vol 4(4), Page 1-11.
- Joseph OS, Jude EO, Joseph OT. (2018). Effect of ethanol stem extract of *Homalium letestuion* gentamicin-induced kidney Injury in rat. Vol. 4(2). Advanced Herbal Medicine. Page 51-64.
- Joseph OT, Olorunfemi AF., Sabastine AZ, Sebastine BS, Joseph OS. (2022). Understanding the cumulative distribution, implication and progress on Covid -19 pandemic as at 7th of February 2022 across different countries of the world: An update report. International Journal of Research and Innovation in Social Science (IJRISS) |Volume VI, Issue IV. Page 691-699.
- Joseph OS, Sabastine AZ, Joseph OT. (2021). Global Implication of Differential Impacts of Covid-19 on Different Countries Using the USA as A Comparism Factor. Journal of Nursing and Health Science. Volume 10, Issue 5. PP 36-44.
- Joseph OS, Builders M, Emem EU, Joseph OT. (2019). Effect of ethanol leaf extract of cassia *angustifolia* extract on liver of wister rats. Global Scientific Journal. Volume 8, Issue 9. Page 1112-11120.
- Joseph OS, Builders M, Emem EU, Joseph OT. (2019). Effect of ethanol leaf extract of *Cassia angustifolia* extract on kidney of Wister Rats. Global Scientific Journal. Volume 8, Issue 9. Page 1023-1031.
- Joseph O.S., Builders M., Joseph O, T., Zubairu S. A., Musa T. (2019). Sub-Acute Toxicity Study of Ethanol Leaf Extract of *Ocimum canum* on Liver of Wister Rats. International Journal of Research and Scientific Innovation. Volume VI (V). Pp. 364-369.
- Joseph OT, Joseph OS, Chinwe AF. (2021). Clinical Study on the Effect of *Moringa oleifera* on Body mass index, Serum Level of High density lipoprotein and Triglyceride in Subjects Taken Tenofovir, Lamivudine and Efavirenz Combination Regimen. J RNA Genom 2021 Volume S04 Issue 004. Page 1-6.
- Joseph OS, Builders M , Joseph OT , Famojuro TI, Ogira JO, Moses FD, Musa TL. (2021). Effect of the Demographic of Covid-19 on Different Countries; Using the USA for Comparism. International journal of multidisciplinary research and analysis. Volume 04 Issue 02. Page 193-203.
- Joseph OS, Builders M, Isinkaye DR, Sebastine AZ, Musa T, Oyepata JP, Joseph OT, Wazis C. (2019). Sub-Acute Toxicity Study of Ethanol Leaf Extract of *Terminalia chebula* On Brain, Stomach and Spleen of Wister Rats. American Journal of

- Biomedical Science & Research. 3(3). Page 277-282.
- Joseph OS, Builders M, Joseph OT, Ariaahu EC, Zubairu SA, Musa T, Joseph OP. (2019). Toxicity study of ethanol leaf extract of *ocimum canum* on heart and lipid profile of wister rats. International Journal of Current Advanced Research. Volume 8. (Issue 05). Page 18800 – 18803.
- Joseph OS, Builders M, Joseph OT, Sabastine AZ, Musa TI, Oyepata PJ. (2019). Sub-acute toxicity study of ethanol leaf extract of *Ocimum canum* on the kidney of wistar rats. African Journal of Pharmaceutical Research & Development. Vol. 11 No.1. Page 1-7.
- Joseph OS, Builders M, Joseph OT, Sabastine AZ, MUSA TL, Oyepata PJ. (2019). Sub-acute toxicity study of ethanol leaf extract of *Ocimum canum* on brain, lungs, stomach and spleen of wister rats. African Journal of Pharmaceutical Research & Development. Vol. 11 No.1. Page 35-42.
- Joseph SO, Opeyemi JT. (2021). Effect of Clinical Study of *Moringa oleifera* on Body mass index, Low density lipoprotein and Triglyceride level in Patients on Tenofovir/lamivudine/efavirenz Combination Therapy. Advanced Herbal Med. Vol. 6. Issue 1. Page. 14-27
- Kluwe WM (1981). Reanl functions tests as indicators of kidney injury in subacute toxicity studies. Toxicol. Appl. Pharmacol. 57:414–424.
- Modupe IB, SOyepata SJ, Akpobome RV (2019). Effect of *Parkia biglobosa* extract on open skin wound healing in dexamethasone - induced hyperglycaemia and histological assessment in rats. African Journal of Pharmacy and Pharmacology. Vol. 13(8), pp. 84-89.
- Okokon JE, Joseph OS, Emem EE. (2016). Hepatoprotective activity of *Homalium letestui* stem extract against paracetamol liver injury. Avicenna Journal of Phytomedicine. 13(4): 87 – 92.
- Okokon JE, Joseph OS, Emem EE. (2016). Nephroprotective activity of *Homalium letestui* stem extract against paracetamol induced kidney injury. Journal of Experimental and Integrative Medicine. Volume 6 (1): 38-43.
- Oluwakanyesola AS, Joseph OS., Jacob A, Rebecca SM, Joseph OT. (2018). Sub-acute haematological toxicity study of safi® blood purifier on wister rats. The Nigerian Journal of Pharmacy. Volume 52 (20).
- Oyebadejo SA, Joseph OS, Adesite SO, Omorilewa AO. (2019). Effect of Citrus Limon Juice and Tamoxifen on the Tumour growth mass Indices, Cell Proliferation, Cell Viability and Cytogenetic (Mitotic Index) of Sprague Dawley Rats Induced MCF-7 Breast Cancer Cells. Saudi Journal of Biomedical Research. (4). Pg. 216 - 225.
- Oyepata JS, Simeon JO. (2022). The Earth: An Alien Planet in Another Universe. Global Journal of Science Frontier Research: A Physics and Space Science. Volume 22 Issue 1. Page 55-57.
- Oyepata JS. (2021). The Earth: A Lost Planet from another Universe. International Journal Of Multidisciplinary Research And Analysis. Volume 04 Issue 12. Page 1795-1797

Palani S, Raja S, Rajalingam D, Praveen KR, Senthil, KB. (1997). Therapeutic efficacy of *Pimpinella tirupatiensis* (Apiaceae) on acetaminophen induced nephrotoxicity and oxidative stress in male albino rats. International Journal of PharmTech Research. 1 (3): 925-934.

Rai S, Ghosh H, Basheer M. (1997). Phytochemical Characterization and Antioxidative Property of *Ocimum canum*: Effect of Ethanolic Extract of Leaves and Seeds on Basic Immunologic and Metabolic Status of Male Rats. J Immuno Biol. 2016, 1:108.

Sabastine AZ, Joseph OS, Joseph OS, Famojuro TI, Olorunfemi AF. (2021). Effect of Cashew apple juice (*Anacardium occidentale* L.) on Hematology and Spleen of Gentamicin Induced Injury in Albino Rats. Global Scientific Journal. Volume 9, Issue 7. Page 3686-3698.

Sabastine AZ, Musa TL, Joseph OS, Builders M, Joseph OT. (2019). Histological study of effect of ethanol stem extracts of *Homalium letestui* in paracetamol induced injury in albino rat, using various staining techniques. American Journal of Biomedical Science & Research. 4(2). Page 82 – 89.

Samson AO, Joseph OS, Samson OA, Emem RA. (2019). Effect of Citrus Linton Juice and Tamoxifen on The oxidative activities of MCT-7 cell induced Breast Cancer in Sprague Dawley Rats. Saudi Journal of Biomedical Research. Volume 8 (7). Page 76-92.

Simeon JO, Builders M, Haruna WC, Tosin JO, Zubairu SA, Lubo MT. (2019). Effect of administration ethanol leaf extract of terminaliachebula on liver of wister rat. International Journal of Research and Scientific Innovation. Volume VI (Issue VII). Page 91- 97.

Simeon JO, C Ariahu Emmanuel, Tosin JO, Zubairu SA. (2022). Virological and immunological consequences of Covid -19 pandemic distribution across different countries; A seven days update study. International Journal of Advances in Engineering and Management (IJAEM) Volume 4, Issue 8. pp: 871-883

Simeon JO, Modupe B, Haruna WC, Zubairu SA, Lubo MT, Tosin JO. (2019). Histological study of effect of ethanol stem extracts of *Homalium letestui* on thioacetamide - induced injury in albino rat, using various staining techniques. International Journal of Research and Scientific Innovation. Volume VI (Issue VII). Page 77 – 85.

Simeon JO, Simeon JO, Zubairu SA, Adegbeniga AD (2021). Concomitant administration of ethanol leaf extract of *Thymus vulgaris* on Diazepam– induced Sedation and Hypnosis in Wister Rat. Journal of Nursing and Health Science. Volume 16, Issue 5. PP 04-09.

Simeon JO, Tosin JO, Adegbeniga AD. (2021). The Relative Global Consequences of Cumulative Distribution of Covid-19, Using the USA as Comparison Factor and Cumulative Covid -19 Data of 31st October 2021. International Journal of

Multidisciplinary Research And Analysis. Page 1906 -1917.

Simeon JO, Tosin JO, Zubairu SA, Daniel MF. (2022). Toxicological evaluation of *Lavandula stoechas* on heart and blood of wistar rat. International Journal of Advances in Engineering and Management (IJAEM). Volume 4, Issue 4 Apr 2022, pp: 1233-1241.

Simeon JO, Tosin JO, Zubairu SA, Oyepata JS (2022). Studying the trend and progress on Covid-19 pandemic from 29th January to 4th of February 2022 across different countries of the world. International Journal of Research and Innovation in Social Science (IJRISS) |Volume VI, Issue II. Page 499-505.

Simeon JO, Zubairu SA, Tosin JO (2021). Clinical evaluation of the potential benefits of taking *Moringaoleifera* on blood triglyceride and cholesterol level in patient taking Tenofovir/Lamivudine/Efavirenz (TLE) combination. Journal of Pharmaceutical Science & Research. Vol. 13(10), 623-629.

Simeon JO, Zubairu SA, Tosin JO, Sunday SB. (2022). Update report and analysis on the global trends and progress of Covid -19 pandemic on 18th January, 2022 across different countries of the world. International Journal of Research and Innovation in Applied Science (IJRIAS) |Volume VII, Issue IV. Page 58 -66.

Simeon, J.O., Tosin, J.O., Zubairu, S.A. (2022). Cumulative evaluation of demography and distribution of COVID-19 around the globe: An update report of COVID-19 until 17th February 2022. Int J

Epidemiol Health Sci;3(6): e34. Doi: 10.51757/IJEHS.3.6.2022.251435.

Smith-Hall C, Larsen O, Pouliot MP. (1997). eople, plants and health: a conceptual framework for assessing changes in medicinal plant consumption. J EthnobiolEthnomed. (2012). 8: 43.

Sofowora EA. (1989). Medicinal Plants and Traditional Medicine in Africa 1st edition, Spectrum Books Ltd. Ibadan, Nigera. 34-40

Solomon IP, Oyebadejo SA, Ukpo EM. Joseph OS. (2015). Changes in serum electrolyte, creatinine and urea of fresh Citrus limon juice administered to growing rabbits (*Oryctolagus cuniculus*). International Journal of Agricultural Science Research. Vol. 4(8), pp. 180-183.

Solomon IP, Oyebadejo SA, Ukpo EM. Joseph OS. (2015). Effect of Fresh Citrus limon Juice on Liver Histomorphology of Growing Rabbits (*Oryctolagus cuniculus*). Scholars Journal of Agriculture and Veterinary Sciences.2 (5):347-351.

Timothy S.Y., Wazis C.H., Midala T.A. S, Joseph O.S., Sabastine A.Z., Nachanaa T. and Oiza F.D. (2017). Evaluation of Anti-Diarrhoeal Activity of Different Bark Extracts of *Faidherbiaalbida* (Delile) A (Chav) in Albino Rats. Bima Journal of Science and Technology Vol. 1 (2). Pg. 122-130

Tosin JO, Simeon JO. (2022). Mathematical and demographic understanding on the effect Covid 19 across the country of the world; An update report of cases and death from 2nd to

8th of August, 2022. International Journal of Advances in Engineering and Management (IJAEM) Volume 4, Issue 8. pp: 891-903.

Tosin JO, Wolfe OA, Iyeopu SM, Simeon JO, Chinwe A, Lubo MT. (2019). Clinical study on the effect of *Moringaoleifera* on serum level of glucose and tryglyceride in subjects taken tenofovir, lamivudine and efavirenz combination regimen. European Scientific Journal. Vol.15, (.21). Page 280 -293.

Tosin JO, Zubairu SA, Simeon JO. (2021). Clinical Effect of *Moringaoleifera* on Body Mass Index, Triglyceride and High Density Lipoprotein in Subjects Taken Tenofovir Combination Regimen. European Journal of Biology and Medical Science Research. Vol.9, No.4, pp.6-19.

Wazis CH, Joseph OS, Modupe B, Joseph OP (2020). Effect of Ethanol Leaf Extract of *Pterocarpussantalinus* Extract on Kidney of Wister Rats. Nigerian Biomedical Science Journal Vol. 17 No 1. Page 35-47.

Zubairu SA, Festus OA, Simeon JO, Irabor I, Tosin JO. (2021). Effect of *Anacardiumoccidentale* Fruit Juice Extract on Haematological Parameters and Spleen of Paracetamol Induced Injury in Albino Rats. Global Scientific Journal. Volume 9, Issue 7. Page 1640-1654.

Zubairu SA, Simeon JO, Tosin JO (2021). Effect of ethanol leaf extract of *Terminalia chebula* extract on kidney of wister rats. Global scientific Journal. Volume 9, Issue 2. Page 514-526.

Zubairu SA, Simeon JO, Tosin JO (2022). Analysis and understanding the progress, trend and consequences of Covid -19 pandemic over a seven days period across different countries of the world. International Journal of Advances in Engineering and Management (IJAEM). Volume 4, Issue 2 pp: 1588-1598.